

6. Literatur

- Ahern TJ, Casal JI, Petsko GA & Klibanov AM (1987)
Control of oligomeric enzyme thermostability by protein engineering.
Proc Natl Acad Sci USA **84**, 675-679
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W & Lipman DJ (1997)
Gapped BLAST and PSI-BLAST: a new generation of protein database search programs.
Nucl Acids Res **25**, 3389-3402
- Anderson A & Cooper RA (1970)
Genetic mapping of a locus for triosephosphate isomerase on the genome of *Escherichia coli* K12.
J Gen Microbiol **62**, 329-334
- Argos P, Rossman MG, Grau UM, Zuber H, Frank G & Tratschin JD (1979)
Thermal stability and protein structure.
Biochemistry **18**, 5698-5703
- Balch WE, Fox GE, Magrum LJ, Woese CR & Wolfe RS (1979)
Methanogens: reevaluation of a unique biological group.
Microbiol Rev **43**, 260-296
- Birnboim HC & Doly J (1979)
A rapid alkaline extraction procedure for screening recombinant plasmid DNA.
Nucl Acids Res **7**, 1513-1523
- Biro J, Fabry S, Dietmaier W, Bogedain C & Hensel R (1990)
Engineering thermostability in archaebacterial glyceraldehyde-3-phosphate dehydrogenase.
Hints for the important role of interdomain contacts in stabilizing protein conformation.
FEBS Lett **275**, 130-134
- Blacklow SC, Raines RT, Lim WA, Zamore PD & Knowles JR (1988)
Triosephosphate isomerase catalysis is diffusion controlled.
Biochemistry **27**, 1158-1167
- Blattner FR, Plunkett III G, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA, Rose DJ, Mau B & Shao Y (1997)
The complete genome sequence of *Escherichia coli* K12.
Science **277**, 1453-1462
- Bledig SA, Ramseier TM & Saier MH Jr (1996)
FruR mediates catabolite activation of pyruvate kinase (*pykF*) gene expression in *Escherichia coli*.
J Bacteriol **178**, 280-283
- Borchert TV, Abagyan R, Jänicke R & Wierenga, RK (1994)
Design, creation, and characterization of a stable, monomeric triosephosphate isomerase.
Proc Natl Acad Sci USA **91**, 1515-1518

- Bradford M.M. (1976)
A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye-binding.
Analytical Biochemistry **150**, 61-66
- Branny P, de la Torre F & Garel J-F (1996)
The genes for phosphofructokinase and pyruvate kinase of *Lactobacillus delbrueckii* subsp. *bulgaricus* constitute an operon.
J Bacteriol **178**, 4727-4730
- Brock TD, Brock KM, Belley RT & Weiss RL (1972)
Sulfolobus: A new genus of sulfur-oxidizing bacteria living at low pH and high temperature.
Arch Microbiol **84**, 54-68
- Brown JR, Daar IO, Krug JR & Maquat LE (1985)
Characterization of the functional gene and several processed pseudogenes in the human triosephosphate isomerase gene family.
Mol Cell Biol **5**, 1694-1706
- Brunner NA, Brinkmann H, Siebers B & Hensel R (1998)
NAD⁺-dependent glyceraldehyde-3-phosphate dehydrogenase from *Thermoproteus tenax*: The first identified member of the aldehyde dehydrogenase superfamily is a glycolytic enzyme with unusual regulatory properties.
J Biol Chem **273**, 6149-6156
- Brunner NA (1998)
Physiologische Funktion und phylogenetische Zuordnung zweier Glycerinaldehyd-3-phosphat-Dehydrogenasen des hyperthermophilen Archaeums *Thermoproteus tenax*.
Dissertation, Universität-GH Essen
- Bullock WO, Fernandez JM & Short JM (1987)
XL1-Blue: A high efficiency plasmid transforming *recA Escherichia coli* strain with β-galactosidase selection.
BioTechniques **5**, 376-379
- Bult CJ, White O, Olsen GJ, Zhou L, Fleischmann RD, Sutton GG, Blake JA, Fitzgerald LM, Clayton RA, Gocayne JD, Kervelage AR, Dougherty BA, Tomb J-F, Adams MD, Reich CI, Overbeek R, Kirkness EF, Weinstock KG, Merrick JM, Glodek A, Scott JL, Geoghegan NSM, Weidman JF, Fuhrmann JL, Nguyen D, Utterback TR, Kelley JM, Peterson JD, Sadow PW, Hanna MC, Cotton MD, Roberts KM, Hurst MA, Kaine BP, Borodovsky M, Klenk H-P, Fraser CM, Smith HO, Woese CR & Venter JC (1996)
Complete genome sequence of the methanogenic archaeon, *Methanococcus jannaschii*.
Science **273**, 1058-1073
- Carninci P, Nishiyama Y, Westover A, Itoh M, Nagaoka S, Sasaki N, Okazaki Y, Muramatsu M & Hayashizaki Y (1998)
Thermostabilization and thermoactivation of thermolabile enzymes by trehalose and its application for the synthesis of full length cDNA.
Proc Natl Acad Sci USA **95**, 520-524

Charlier D, Roovers M, Thia-Toong TL, Durbecq V & Glansdorff N (1997)
Cloning and identification of the *Sulfolobus solfataricus lrp* gene encoding an archaeal homologue of the eubacterial leucine-responsive global transcriptional regulator Lrp.
Gene **201**, 63-68

Chevalier C, Saillard C & Bové JM (1990)
Organization and nucleotide sequences of the *Spiroplasma citri* genes for ribosomal protein S2, elongation factor Ts, spiralin, phosphofructokinase, pyruvate kinase and an unidentified protein.
J Bacteriol **172**, 2693-2703.

Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV, Eiglmeier K, Gas S, Barry III CE, Tekaia F, Badcock K, Basham D, Brown D, Chillingworth T, Connor R, Davies R, Devlin K, Feltwell T, Gentles S, Hamlin N, Holroyd S, Hornsby T, Jagels K, Krogh A, McLean J, Moule S, Murphy L, Oliver S, Osborne J, Quail MA, Rajandream MA, Rogers J, Rutter S, Seeger K, Skelton S, Squares S, Squares R, Sulston JE, Taylor K, Whitehead S & Barrell BG (1998)

Deciphering the biology of *Mycobacterium tuberculosis* from the complete genome sequence.
Nature **393**, 537-544

Danson MJ (1993)
Central metabolism of the Archaea
In Kates M, Kushner DJ & Matheson AT (eds.): The biochemistry of Archaea.
Elsevier, Amsterdam, pp 1-24

Deckert G, Warren PV, Gaasterland T, Young WG, Lenox AL, Graham DE, Overbeek R, Snead MA, Keller M, Aujay M, Huber R, Feldman RA, Short JM, Olsen GJ & Swanson RV (1998)

The complete genome of the hyperthermophilic bacterium *Aquifex aeolicus*.
Nature **392**, 353-358

Delboni LF, Mande SC, Rentier-Delrue F, Mainfroid V, Turley S, Vellieux FM, Martial JA & Hol WG (1995)
Crystal structure of recombinant triosephosphate isomerase from *Bacillus stearothermophilus*. An analysis of potential thermostability factors in six isomerases with known three-dimensional structures points to the importance of hydrophobic interactions.
Protein Sci **4**, 2594-2604

de Vos WM, Kengen SWM, Voorhorst WGB & van der Oost J (1998)
Sugar utilization and its control in hyperthermophiles.
Extremophiles **2**, 201-205

Drews G (1983)
Mikrobiologisches Praktikum.
Springer-Verlag, Heidelberg

Eckerskorn C, Mewes W, Goretzki H & Lottspeich F (1988)
A new siliconized glass fiber as support for chemical analysis of electroblotted proteins.
Eur J Biochem **176**, 509-519

- Engler-Blum G, Meier M, Frank J & Müller GA (1993)
Reduction of background problems in nonradioactive Northern and Southern blot analyses enables higher sensitivity than ^{32}P -based hybridizations.
Anal Biochem **210**, 235-244
- Farber GK & Petsko GA (1990)
The evolution of α/β barrel enzymes.
TIBS **15**, 228-234
- Fischer F, Zillig W, Stetter KO & Schreiber G (1983)
Chemolithoautotrophic metabolism of anaerobic extremely thermophilic Archaeabacteria.
Nature **301**, 511-513
- Fitz-Gibbon S, Choi AJ, Miller JH, Stetter KO, Simon MI, Swanson R & Kim UJ (1997)
A fosmid-based genomic map and identification of 474 genes of the hyperthermophilic archaeon *Pyrobaculum aerophilum*.
Extremophiles **1**, 36-51
- Fothergill-Gilmore LA & Michels PAM (1993)
Evolution of glycolysis.
Prog Biophys & Mol Biol **59**, 105-235
- Franzetti B, Sohlberg B, Zaccai G & von Gabain A (1997)
Biochemical and serological evidence for an RNase E-like activity in halophilic Archaea.
J Bacteriol **179**, 1180-1185
- Fraser CM, Gocayne JD, White O, Adams MD, Clayton RA, Fleischmann RD, Bult CJ, Kerlavage AR, Sutton G, Kelley JM, Fritschman JL, Weidman JF, Small KV, Sansusky M, Fuhrmann J, Nguyen D, Utterback TR, Saudek DM, Phillips CA, Merrick JM, Tomb J-F, Dougherty BA, Bott KF, Hu P-C, Lucier TS, Peterson SN, Smith HO, Hutchinson III CA & Venter JC (1995)
The minimal gene complement of *Mycoplasma genitalium*.
Science **270**, 397-403
- Fürste JP, Pansegrouw W, Frank R, Blöcker H, Scholz P, Bagdasarian M & Lanka E (1986)
Molecular cloning of the plasmid RP4 primase region in a multi-host-range *tacP* expression vector.
Gene **48**, 119-131
- Gotzes A (1996)
Strukturuntersuchungen an der Triose-3-phosphat-Isomerase von *Methanothermus fervidus*.
Examensarbeit, Universität-GH Essen
- Grosjean H & Fiers W (1982)
Preferential codon usage in prokaryotic genes: the optimal codon-anticodon interaction energy and the selective codon usage in efficiently expressed genes.
Gene **18**, 199-209

Hain J, Reiter WD, Hüdepohl U & Zillig W (1992)
Elements of an archaeal promoter defined by mutational analysis.
Nucl Acids Res **20**, 5423-5428

Hanahan D (1983)
Studies on transformation of *Escherichia coli* with plasmids.
J Mol Biol **166**, 557-580

Hennig M, Sterner R, Kirschner K & Jansonius JN (1997)
Crystal structure at 2.0 Å resolution of phosphoribosyl anthranilate isomerase from the hyperthermophile *Thermotoga maritima*: possible determinants of protein stability.
Biochemistry **36**, 6009-6016

Hensel R, Laumann S, Lang J, Heumann H & Lottspeich F (1987)
Characterization of two D-glyceraldehyde-3-phosphate dehydrogenases from the extremely thermophilic archaebacterium *Thermoproteus tenax*.
Eur J Biochem **170**, 325-333

Hensel R & König H (1988)
Thermoadaptation of methanogenic bacteria by intracellular ion concentration.
FEMS Microbiol Letters **49**, 75-79

Hensel R, Schramm A, Hess D & Russell RJM (1998)
3-phosphoglycerate kinase and triosephosphate isomerase from hyperthermophilic archaea: features of biochemical thermoadaptation.
In: Wiegand and Adams (editors), Thermophiles: the key to molecular evolution and the origin of life?

Heppner P (1990)
Klonierung und Sequenzierung des 3-Phosphoglycerat-Kinase-Gens aus dem mesophilen Archaeabakterium *Methanobacterium bryantii*.
Diplomarbeit, Ludwig-Maximilians-Universität München

Hess D, Krüger K, Knappik A, Palm P & Hensel, R (1995)
Dimeric 3-phosphoglycerate kinases from hyperthermophilic Archaea. Cloning, sequencing and expression of the 3-phosphoglycerate kinase gene of *Pyrococcus woesei* in *Escherichia coli* and characterization of the protein. Structural and functional comparison with the 3-phosphoglycerate kinase of *Methanothermus fervidus*.
Eur J Biochem **233**, 227-237

Heukeshoven J & Dernick R (1985)
Simplified method for silver staining of proteins in polyacrylamide gels and the mechanism of silver staining.
Electrophoresis **6**, 103-112

Hill R (1925)
A measure of cooperativity.
Proc R Soc **B100**, 419

- Horlacher R, Xavier KB, Santos H, DiRuggiero J, Kossman M & Boos W (1998)
Archaeal binding protein-dependent ABC transporter: molecular and biochemical analysis of the trehalose/maltose transport system of the hyperthermophilic archaeon *Thermococcus litoralis*.
J Bacteriol **180**, 680-689
- Jetten MS, Gubler ME, Lee SH & Sinskey AJ (1994)
Structural and functional analysis of pyruvate kinase from *Corynebacterium glutamicum*.
Appl Environ Microbiol **60**, 2501-2507
- Jin S & Sonenshein AL (1994)
Identification of two distinct *Bacillus subtilis* citrate synthase genes.
J Bacteriol **176**, 4669-4679
- Jungblut P, Eckerskorn C, Lottspeich F & Klose J (1990)
Blotting efficiency investigated by using two-dimensional electrophoresis, hydrophobic membranes and proteins from different sources.
Electrophoresis **11**, 581-588
- Jurica MS, Mesecar A, Heath PJ, Shi W, Nowak N & Stoddard BL (1998)
The allosteric regulation of pyruvate kinase by fructose-1,6-bisphosphate.
Structure **6**, 195-210
- Kawarabayasi Y, Sawada M, Horikawa H, Haikawa Y, Hino Y, Yamamoto S, Sekine M, Baba S, Kosugi H, Hosoyama A, Nagai Y, Sakai M, Ogura K, Otuka R, Nakazawa H, Takamiya M, Ohfuku Y, Funahashi T, Tanaka T, Kudoh Y, Yamazaki J, Kushida N, Oguchi A, Aoki K, Nakamura Y, Robb TF, Horikoshi K, Masuchi Y, Shizuya H & Kikuchi H (1998)
Complete sequence and gene organization of the genome of a hyperthermophilic Archaeabacterium, *Pyrococcus horikoshii* OT3.
DNA Research **5**, 55-76
- Keeling PJ & Doolittle WF (1997)
Evidence that eukaryotic triosephosphate isomerase is of alpha-proteobacterial origin.
Proc Natl Acad Sci USA 1270-1275
- Kengen SWM, de Bok FAM, van Loo N-D, Dijkema C, Stams AJM & de Vos WM (1994)
Evidence for the operation of a novel Embden-Meyerhof-pathway that involves ADP-dependent kinases during sugar fermentation by *Pyrococcus furiosus*.
J Biol Chem **269**, 17537-17541
- Kengen SWM, Tuininga JE, de Bok FAM, Stams AJM & de Vos WM (1995)
Purification and characterization of a novel ADP-dependent glucokinase from the hyperthermophilic archaeon *Pyrococcus furiosus*.
J Biol Chem **270**, 30453-30475

Klenk H-P, Clayton RA, Tomb J-F, White O, Nelson KE, Ketchum KA, Dodson RJ, Gwinn M, Hickey EK, Peterson JD, Richardson DL, Kerlavage AR, Graham DE, Kyrpides NC, Fleischmann RD, Quackenbush J, Lee NH, Sutton GG, Gill S, Kirkness EF, Dougherty BA, McKenney K, Adams MD, Loftus B, Scott P, Reich CI, McNeil LK, Badger JH, Glodek A, Zhou L, Overbeek R, Gocayne JD, Weidman JF, McDonald L, Utterback T, Cotton MD, Spriggs T, Artiach P, Kaine BP, Sykes SM, Sadow PW, D'Andrea KP, Bowman C, Fujii C, Garland SA, Mason TM, Olsen GJ, Fraser CM, Smith HO, Woese CR & Venter JC (1997)
The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*.
Nature **390**, 364-370

Kletzin A, Mukund S, Kelley-Crouse TL, Chan MK, Rees DC & Adams MW (1995)
Molecular characterization of the genes encoding the tungsten-containing aldehyde ferredoxin oxidoreductase from *Pyrococcus furiosus* and formaldehyde ferredoxin oxidoreductase from *Thermococcus litoralis*.
J Bacteriol **177**, 4817-4819

Knowles, JR (1991)
Enzyme catalysis: Not different, just better.
Nature **350**, 121-124

Kohl L, Callens M, Wierenga RK, Opperdoes F & Michels PA (1994)
Triose-phosphate isomerase of *Leishmania mexicana mexicana*: Cloning and characterization of the gene, overexpression in *Escherichia coli* and analysis of the protein.
Eur J Biochem **220**, 331-338

Kohlhoff M (1995)
Struktur und Funktion der Triose-3-phosphat-Isomerase bei hyperthermophilen Archaea.
Dissertation, Universität-GH Essen

Kohlhoff M, Dahm A & Hensel R (1996)
Tetrameric triosephosphate isomerase from hyperthermophilic Archaea.
FEBS Letters **383**, 245-250

Krüger K, Hermann T, Armbruster V & Pfeifer F (1998)
The transcriptional activator GvpE for the halobacterial gas vesicle genes resembles a basic region leucine-zipper regulatory protein.
J Mol Biol **279**, 761-771

Kuo YP, Thompson DK, St Jean A, Charlebois RL & Daniels CJ (1997)
Characterization of two heat shock genes from *Haloferax volcanii*: a model system for transcription regulation in the Archaea.
J Bacteriol **179**, 6318-6324

Kuroda S, Otaka S & Fujisawa Y (1994)
Fermentable and nonfermentable carbon sources sustain constitutive levels of expression of yeast triosephosphate dehydrogenase 3 gene from distinct promoter elements.
J Biol Chem **269**, 6153-62

- Laemmli UK (1970)
Cleavage of structural proteins during the assembly of the head of bacteriophage T4.
Nature **277**, 680-685
- Lamosa P, Martins LO, da Costa MS & Santos H (1998)
Effects of temperature, salinity, and medium composition on compatible solute accumulation
by *Thermococcus* spp..
Appl Environ Microbiol **64**, 3591-3598
- Langley KE, Villarejo MR, Fowler AV, Zamenhof PJ & Zabin I (1975)
Molecular basis of β -galactosidase α -complementation.
Proc Natl Acad Sci USA **72**, 1254-1257
- Larsen TM, Laughlin LT, Holden HM, Rayment I & Reed GH (1994)
Structure of rabbit muscle pyruvate kinase complexed with Mn^{2+} , K^+ , and pyruvate.
Biochemistry **33**, 6301-6309
- Larsen TM, Benning MM, Rayment I & Reed GH (1998)
Structure of the bis(Mg^{2+})-ATP-oxalate complex of the rabbit muscle pyruvate kinase at 2.1 Å
resolution: ATP binding over a barrel.
Biochemistry **37**, 6247-6255
- Leinfelder W, Jarsch M & Böck A (1985)
The phylogenetic position of the sulfur-dependent Archaeabacterium *Thermoproteus tenax*:
Sequence of its 16S rRNA gene.
System Appl Microbiol **6**, 164-170
- Leyva-Vazquez MA & Setlow P (1994)
Cloning and nucleotide sequence of the genes encoding triosephosphate isomerase,
phosphoglycerate mutase and enolase from *Bacillus subtilis*.
J Mol Biol **176**, 3903-3910
- Lineweaver H & Burk D (1934)
The determination of enzyme dissociation constants.
J Am Chem Soc **56**, 658-660
- Lolis E & Petsko GA (1990)
Crystallographic analysis of the complex between triosephosphate isomerase and
2-phosphoglycolate at 2.5 Å: Implications for catalysis.
Biochemistry **29**, 6619-6625
- Macedo-Ribeiro S, Darimont B, Sterner R, Huber R (1996)
Small structural changes account for the high thermostability of 1[4Fe-4S] ferredoxin from
the hyperthermophilic bacterium *Thermotoga maritima*.
Structure **4**, 1291-1301
- Maitra PK & Lobo Z (1971)
Control of glycolytic enzyme synthesis in yeast by products of the hexokinase reaction.
J Biol Chem **246**, 489-499

- Makrides SC (1996)
Strategies for achieving high-level expression of genes in *Escherichia coli*.
Microbiol Rev **60**, 512-538
- Martins LO, Huber R, Huber K, Stetter KO, da Costa M & Santos H (1997)
Organic solutes in hyperthermophilic archaea.
Appl Env Microbiol **63**, 896-902
- Mattevi A, Valentini G, Rizzi M, Speranza ML, Bolognesi M & Coda A (1995)
Crystal structure of *Escherichia coli* pyruvate kinase type I: molecular basis of the allosteric transition.
Structure **3**, 729-41
- Matussek K, Moritz P, Brunner N, Eckerskorn C, Hensel R (1998)
Cloning, sequencing, and expression of the gene encoding cyclic 2, 3-diphosphoglycerate synthetase, the key enzyme of cyclic 2,3-diphosphoglycerate metabolism in *Methanothermus fervidus*.
J Bacteriol **180**, 5997-6004
- Meakin SA, Nash J, Murray WD, Kennedy KJ & Sprott GD (1991)
A generally applicable technique for the extraction of restrictionable DNA from methanogenic bacteria.
J Microbiol Meth **14**, 119-126
- Meijer WG, de Boer P & van Keulen G (1997)
Xanthobacter flavus employs a single triosephosphate isomerase for heterotrophic and autotrophic metabolism.
Microbiology **143**, 1925-1931
- Menendez-Arias L & Argos P (1989)
Engineering protein thermal stability. Sequence statistics point to residue substitutions in α -helices.
J Mol Biol **206**, 397-406
- Muir JM, Russell RJ, Hough DW & Danson MJ (1995)
Citrate synthase from the hyperthermophilic Archaeon, *Pyrococcus furiosus*.
Protein Eng **8**, 583-592
- Muirhead H, Clayden DA, Barford D, Lorimer CG, Fothergill-Gilmore LA, Schiltz E & Schmitt W (1986)
The structure of cat muscle pyruvate kinase.
EMBO J **5**, 475-481
- Mukund S & Adams MMW (1995)
Glyceraldehyde-3-phosphate ferredoxin oxidoreductase, a novel tungsten-containing enzyme with a potential glycolytic role in the hyperthermophilic archaeon *Pyrococcus furiosus*.
J Biol Chem **270**, 8389-8392
- Mullis KB, Falloona S, Scharf R, Saiki RK, Horn G & Ehrlich H (1986)
Specific enzymatic amplification of DNA *in vitro*: the polymerase chain reaction.
Cold Spring Harbor Symposium of Quantitative Biology **51**, 263-273

Nairn J, Smith S, Allison PJ, Rigden D, Fothergill-Gilmore LA & Price NC (1995)
Cloning and sequencing of a gene encoding pyruvate kinase from *Schizosaccharomyces pombe*; implications for quaternary structure and regulation of the enzyme.
FEMS Microbiol Lett **134**, 221-226

Nairn J, Duncan D, Gray LM, Urquhart G, Binnie M, Byron O, Fothergill-Gilmore LA & Price NC (1998)
Purification and characterization of pyruvate kinase from *Schizosaccharomyces pombe*: Evidence for an unusual quaternary structure.
Protein Expr Purif **14**, 247-253

Pawluk A, Scopes RK & Griffiths-Smith K (1986)
Isolation and properties of the glycolytic enzymes from *Zymomonas mobilis*. The five enzymes from glyceraldehyde-3-phosphate dehydrogenase through to pyruvate kinase.
Biochem J **238**, 275-281

Pearson WR & Lipman DJ (1988)
Improved tools for biological sequence comparison.
Proc Natl Acad Sci USA **85**, 2444-2448

Plaut B & Knowles JR (1972)
pH-dependence of the triose-3-phosphate isomerase reaction.
Biochem J **129**, 301-310

Potter S & Fothergill-Gilmore LA (1992)
Purification and properties of pyruvate kinase from *Thermoplasma acidophilum*.
FEMS Microbiol Lett **94**, 235-240

Prodromou C, Artymiuk PJ & Guest JR (1992)
The aconitase of *Escherichia coli*. Nucleotide sequence of the aconitase gene and amino acid sequence similarity with mitochondrial aconitases, the iron-responsive-element-binding protein and isopropylmalate isomerases.
Eur J Biochem **204**, 599-609

Ramos A, Raven NDH, Sharp RJ, Bartolucci S, Rossi M, Cannio R, Lebbink J, van der Oost J, de Vos WM & Santos H (1997)
Stabilization of enzymes against thermal stress and freeze-drying by mannosylglycerate.
Appl Environ Microbiol **63**, 4020-4025

Ramseier TM, Bledig S, Michotey V, Feghali R & Saier MH Jr (1995)
The global regulatory protein FruR modulates the direction of carbon flow in *Escherichia coli*.
Mol Microbiol **16**, 1157-1169

Ranie J, Kumar VP & Balaram H (1993)
Cloning of the triosephosphate isomerase gene of *Plasmodium falciparum* and expression in *Escherichia coli*.
Mol Biochem Parasitol **61**, 159-169

- Rentier-Delrue F, Mande SC, Moyens S, Terpstra P, Mainfroid V, Goraj K, Lion M, Hol WG & Martial JA (1993)
Cloning and overexpression of the triosephosphate isomerase genes from psychrophilic and thermophilic bacteria. Structural comparison of the predicted protein sequences.
J Mol Biol **229**, 85-93
- Russell RJM, Campbell-Ferguson JM, Hough DW, Danson MJ & Taylor GL (1997)
Determinants of protein hyperthermostability revealed by the crystal structure of *Pyrococcus furiosus* citrate synthase.
Biochemistry **36**, 9983-9994
- Russell RJM, Gerike U, Danson MJ, Hough DW & Taylor GL (1998)
Structural adaptations of the cold-active citrate synthase from an Antarctic bacterium.
Structure **6**, 351-361
- Sabnis NA, Yang H & Romeo T (1995)
Pleiotropic regulation of central carbohydrate metabolism in *Escherichia coli* via the gene *csrA*.
J Biol Chem **270**, 29096-29104
- Saiki RK, Gelfand DH, Stoffel S, Scharf S, Higuchi RH, Horn GT, Mullis KB & Ehrlich HA (1988)
Primer-directed enzymatic amplification of DNA with a thermostable polymerase.
Science **239**, 487-490
- Sakai H & Ohta T (1987)
Evidence for two activated forms of pyruvate kinase from *Bacillus stearothermophilus* in the presence of ribose-5-phosphate.
J Biochem (Tokyo) **101**, 633-642
- Sambrook J, Fritsch EF & Maniatis T (1989)
Molecular cloning. A laboratory manual.
Cold Spring Harbor Laboratory Press, New York
- Sanger F., Nicklen S. & Coulson A.R. (1977)
DNA-Sequencing with chain-terminating inhibitors
Proc Natl Acad Sci USA **74**, 5463-5467
- Schäfer S, Götz M, Eisenreich W, Bacher A & Fuchs G (1989)
¹³C-NMR study of autotrophic CO₂ fixation in *Thermoproteus neutrophilus*.
Eur J Biochem **184**, 151-156
- Schägger H & von Jagow G (1987)
Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kD.
Anal Biochem **166**, 368-379
- Schleper C, de Long EF, Preston CM, Feldman RA, Wu K-Y & Swanson RV (1998)
Genomic analysis reveals chromosomal variation in natural populations of the uncultured psychrophilic archaeon *Cenarchaeum symbiosum*.
J Bacteriol **180**, 5003-5009

- Schliebs W, Thanki N, Eritja R & Wierenga R (1996)
Active site properties of monomeric triosephosphate isomerase (monoTIM) as deduced from mutational and structural studies.
Prot Sci **5**, 229-239
- Schmidt M, Svendsen I & Feierabend J (1995)
Analysis of the primary structure of the chloroplast isozyme of triosephosphate isomerase from rye leaves by protein and cDNA sequencing indicates a eukaryotic origin of its gene.
Biochim Biophys Acta **1261**, 257-264
- Scholz S, Sonnenbichler J, Schäfer W & Hensel R (1992)
Di-*myo*-inositol-1,1'-phosphate: a new inositol phosphate isolated from *Pyrococcus woesei*.
FEBS Lett **306**, 239-242
- Scholz S, Wolff S & Hensel R (1998)
The biosynthesis pathway of Di-*myo*-inositol-1,1'-phosphate in *Pyrococcus woesei*.
FEMS Microbiol Lett **168**, 37-42
- Schurig H, Beaucamp N, Ostendorp R, Jaenicke R, Adler E & Knowles JR (1995)
Phosphoglycerate kinase and triosephosphate isomerase from the hyperthermophilic bacterium *Thermotoga maritima* form a covalent bifunctional enzyme complex.
EMBO J **14**, 442-451
- Selig M, Xavier KB, Santos H & Schönheit P (1997)
Comparative analysis of Embden-Meyerhof and Entner-Doudoroff glycolytic pathways in hyperthermophilic archaea and the bacterium *Thermotoga*.
Arch Microbiol **167**, 217-232
- Shaw-Lee RL, Lissemore JL & Sullivan DT (1991)
Structure and expression of the triose phosphate isomerase (*tpi*) gene of *Drosophila melanogaster*.
Mol Gen Genet **230**, 225-229
- Shuman S (1994)
Novel approach to molecular cloning and polynucleotide synthesis using vaccinia DNA topoisomerase.
J Biol Chem **269**, 32678-32684
- Siebers B & Hensel R (1993)
Glucose catabolism of the hyperthermophilic archaeum *Thermoproteus tenax*.
FEMS Microbiol Lett **111**, 1-8
- Siebers B (1995)
Untersuchungen zum Kohlenhydrat-Metabolismus des hyperthermophilen Archaeums *Thermoproteus tenax*.
Dissertation, Universität-GH Essen
- Siebers B, Wendisch VF & Hensel R (1997)
Carbohydrate metabolism in *Thermoproteus tenax*: *in vivo* utilization of the non-phosphorylative Entner-Doudoroff pathway and characterization of its first enzyme, glucose dehydrogenase.
Arch Microbiol **168**, 120-127

Siebers B, Klenk HP & Hensel R (1998)

PPi-dependent phosphofructokinase from *Thermoproteus tenax*, an archaeal descendant of an ancient line in phosphofructokinase evolution.

J Bacteriol **180**, 2137-2143

Smith DR, Doucette-Stamm LA, Deloughery C, Lee H, Dubois J, Aldredge T, Bashirzadeh R, Blakeley D, Cook R, Gilbert K, Harrison D, Hoang L, Keagle P, Lumm W, Pothier B, Qiu D, Spadafora R, Vicaire R, Wang Y, Wierzbowski J, Gibson R, Jiwani N, Caruso A, Bush D, Sfer H, Patwell D, Prabhakar S, McDougall S, Shimer G, Goyal A, Pietrovski S, Church G, Daniels CJ, Mao J-I, Rice P, Nölling J & Reeve JN (1997)

Complete genome sequence of *Methanobacterium thermoautotrophicum* ΔH: Functional analysis and comparative genomics.

J Bacteriol **179**, 7135-7155

Southern EM (1975)

Detection of specific sequences among DNA-fragments separated by gel electrophoresis.

J Mol Biol **98**, 503-517

Sterner R, Kleemann GR, Szadkowski H, Lustig A, Hennig M & Kirschner K (1996)

Phosphoribosyl anthranilate isomerase from *Thermotoga maritima* is an extremely stable and active homodimer.

Protein Sci **5**, 2000-2008

Stetter KO, Thomm M, Winter J, Wildgruber G, Huber H, Zillig W, Janekovic D, König H, Palm P & Wunderl S (1981)

Methanothermus fervidus sp. nov., a novel extremely thermophilic methanogen isolated from an icelandic hot spring.

Zbl Bacteriol Hyg I Abt Orig **C 2**, 166-178

Stetter KO (1988)

Archaeoglobus fulgidus gen. nov., sp. nov.: a new taxon of extremely thermophilic archaeabacteria.

Syst Appl Microbiol **10**, 172-173

Stetter KO (1996)

Hyperthermophilic prokaryotes.

FEMS Microbiol Rev **18**, 149-158

Studier FW & Moffatt BA (1986)

Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes.

J Mol Biol **189**, 113-130

Swinkels BW, Gibson WC, Osinga KA, Kramer R, Veeneman GH, van Boom JH & Borst P (1986)

Characterization of the gene for the microbody (glycosomal) triosephosphate isomerase of *Trypanosoma brucei*.

EMBO J **5**, 1291-1298

Swofford DL (1996)

PAUP*: Phylogenetic analysis using parsimony (and other methods), version 4.0.
Sinauer Associates, Sunderland, Massachusetts

Timasheff SN & Arakawa T (1989)

Stabilization of protein structure by solvents.

In: Creighton (editor), Protein structure, a practical approach, IRL Press.

Thein SL & Wallace RB (1986)

The use of synthetic oligonucleotides as specific hybridization probes in the diagnosis of genetic disorders.

In: Davis (editor), Human genetic diseases, a practical approach, IRL Press.

Thompson JD, Higgins DG & Gibson TJ (1994)

CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice.

Nucl Acids Res **22**, 4673-4680

van der Oost J, Schut G, Kengen SWM, Hagen WR, Thomm M & de Vos WM (1998)

The ferredoxin-dependent conversion of glyceraldehyde-3-phosphate in the hyperthermophilic archaeon *Pyrococcus furiosus* represents a novel site of glycolytic regulation.

J Biol Chem **273**, 28149-28154

Villeret V, Clantin B, Tricot C, Legrain C, Roovers M, Stalon V, Glansdorff N, Van Beeumen J (1998)

The crystal structure of *Pyrococcus furiosus* ornithine carbamoyltransferase reveals a key role for oligomerization in enzyme stability at extremely high temperatures.

Proc Natl Acad Sci USA **95**, 2801-2806

Walter AE, Turner DH, Kim J, Lyttle MH, Müller P, Mathews DH & Zuker M (1994)

Coaxial stacking of helices enhances binding of oligoribonucleotides and improves predictions of RNA folding.

Proc Natl Acad Sci USA **91**, 9218-9222

Waygood EB, Rayman MK & Sanwal BD (1975)

The control of pyruvate kinases of *Escherichia coli*. II. Effectors and regulatory properties of the enzyme activated by ribose 5-phosphate.

Can J Biochem **53**, 444-454

Weil CF, Cram DS, Sherf BA & Reeve JN (1988)

Structure and comparative analysis of the genes encoding components C of methyl coenzyme M reductase in the extremely thermophilic archaeabacterium *Methanothermus fervidus*.

J Bacteriol **70**, 4718-4726

Wierenga RK, Noble MEM & Davenport RC (1992)

Comparison of the refined crystal structures of liganded and unliganded chicken, yeast and trypanosomal triosephosphate isomerase.

J Mol Biol **224**, 1115-1126

Wilson TJG, Bertrand N, Tang JL, Feng JX, Pan MQ, Barber CE, Dow JM & Daniels MJ (1998)

The *rpfA* gene of *Xanthomonas campestris* pathovar *campestris*, which is involved in regulation of pathogenicity factor production, encodes an aconitase.

Mol Microbiol **28**, 961-971

Woese CR, Kandler O & Wheelis ML (1990)

Towards a natural system of organisms: proposal for the domains Archaea, Bacteria, and Eucarya.

Proc Natl Acad Sci USA **87**, 4576-4579

Yanisch-Perron C, Vieira J & Messing J (1985)

Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors.

Gene **33**, 103-119

Yu J-P, Ladapo J & Whitman WB (1994)

Pathway of glycogen metabolism in *Methanococcus maripaludis*.

J Bacteriol **176**, 325-332

Zillig W, Stetter KO, Schäfer W, Janekovic D, Wunerl S, Holz I & Palm P (1981)

Thermoproteales: a novel type of extremely thermoacidophilic anaerobic Archaeabacteria isolated from Icelandic solfatares.

Zentralbl Bakteriol Hyg Abt Orig C **2**, 205-227

Zillig W, Holz I, Klenk HP, Trent J, Wunderl S, Janekovic D, Insel E & Haas B (1987)

Pyrococcus woesei, sp. nov., an ultra-thermophilic marine archaeabacterium, representing a novel order, Thermococcales.

System Appl Microbiol **9**, 62-70

Zwickl P, Fabry S, Bogedain C, Haas A & Hensel R (1990)

Glyceraldehyde-3-phosphate dehydrogenase from the hyperthermophilic archaeabacterium *Pyrococcus woesei*: characterization of the enzyme, cloning and sequencing of the gene, and expression in *Escherichia coli*.

J Bacteriol **172**, 4329-4338