

## Literatur

Angrand,P.O., Woodroffe,C.P., Buchholz,F., and Stewart,A.F. (1998). Inducible expression based on regulated recombination: a single vector strategy for stable expression in cultured cells. *Nucleic Acids Res.* 26, 3263-3269.

Asfari,M., Janjic,D., Meda,P., Li,G., Halban,P.A., and Wollheim,C.B. (1992). Establishment of 2-mercaptoethanol dependent differentiated insulin secreting cell lines. *Endocrinology* 130, 167-178.

Bailly,A., Spaeth,G., Bender,V., and Weiss,M.C. (1998). Phenotypic effects of the forced expression of HNF4 and HNF1 $\alpha$  are conditioned by properties of the recipient cell. *J. Cell Sci.* 111, 2411-2421.

Barbacci,E., Reber,M., Ott,M., Breillat,C., Huetz,F., and Cereghini,S. (1999). Variant hepatocyte nuclear factor 1 is required for visceral endoderm specification. *Development* 126, 4795-4805.

Barroso,I., Gurnell,M., Crowley,V.E., Agostini,M., Schwabe,J.W., Soos,M.A., Maslen,G.L., Williams,T.D., Lewis,H., Schafer,A.J., Chatterjee,V.K., and O'Rahilly,S. (1999). Dominant negative mutations in human PPAR $\gamma$  associated with severe insulin resistance, diabetes mellitus and hypertension. *Nature* 402, 880-883.

Bingham,C., Ellard,S., Allen,L., Bulman,M., Shepherd,M., Frayling,T., Berry,P.J., Clark,P.M., Lindner,T., Bell,G.I., Ryffel,G.U., Nicholls,A.J., and Hattersley,A.T. (2000). Abnormal nephron development associated with a frameshift mutation in the transcription factor hepatocyte nuclear factor 1 beta. *Kidney Int.* 57, 898-907.

Bogan,A.A., Dallas-Yang,Q., Ruse,M.D., Jr., Maeda,Y., Jiang,G., Nepomuceno,L., Scanlan,T.S., Cohen,F.E., and Sladek,F.M. (2000). Analysis of protein dimerization and ligand binding of orphan receptor HNF4 $\alpha$ . *J. Mol. Biol.* 302, 831-851.

Bulla,G.A. (1997). Selective loss of the hepatic phenotype due to the absence of a transcriptional activation pathway. *Somat. Cell Mol. Genet.* 23, 185-201.

Bulman,M.P., Dronsfield,M.J., Frayling,T., Appleton,M., Bain,S.C., Ellard,S., and Hattersley,A.T. (1997). A missense mutation in the hepatocyte nuclear factor 4 $\alpha$  gene in a UK pedigree with maturity onset diabetes of the young. *Diabetologia* 40, 859-862.

Bulman,M.P., Ellard,S., and Hattersley,A.T. (2000). R127W in HNF4 $\alpha$  is a loss of function mutation causing maturity-onset diabetes of the young (MODY) in a UK Caucasian family. *Diabetologia* 43, 1203.

Burris,T.P. (2001). The nuclear receptor superfamily. In: *Nuclear receptors and genetic disease*, T.P.Burris and E.R.B.McCabe (eds.), Academic Press, pp. 1-58.

Carew,J.A., Pollak,E.S., Lopaciuk,S., and Bauer,K.A. (2000). A new mutation in the HNF4 binding region of the factor VII promoter in a patient with severe factor VII deficiency. *Blood* 96, 4370-4372.

- Cereghini,S. (1996). Liver-enriched transcription factors and hepatocyte differentiation. *FASEB J.* 10, 267-282.
- Chen,W.S., Manova,K., Weinstein,D.C., Duncan,S.A., Plump,A.S., Prezioso,V.R., Bachvarova,R.F., and Darnell,J.E., Jr. (1994). Disruption of the HNF4 gene, expressed in visceral endoderm, leads to cell death in embryonic ectoderm and impaired gastrulation of mouse embryos. *Genes Dev.* 8, 2466-2477.
- Chung,A.C.K. and Cooney,A.J. (2001). Retinoid receptors. In: Nuclear receptors and genetic disease, T.P.Burris and E.R.B.McCabe (eds.), Academic press, pp. 245-295.
- Coucouvanis,E. and Martin,G.R. (1999). BMP signaling plays a role in visceral endoderm differentiation and cavitation in the early mouse embryo. *Development* 126, 535-546.
- Dang,C.V. and Semenza,G.L. (1999). Oncogenic alterations of metabolism. *Trends Biochem. Sci.* 24, 68-72.
- Dell,H. and Hadzopoulou-Cladaras,M. (1999). CREB-binding protein is a transcriptional coactivator for hepatocyte nuclear factor 4 and enhances apolipoprotein gene expression. *J. Biol. Chem.* 274, 9013-9021.
- Dressel,U. and Baniahmad,A. (2001). Thyroid hormone receptors. In: Nuclear receptors and genetic disease, T.P.Burris and E.R.B.McCabe (eds.), Academic press, pp. 59-96.
- Drewes,T., Clairmont,A., Klein-Hitpass,L., and Ryffel,G.U. (1994). Estrogen inducible derivatives of hepatocyte nuclear factor 4, hepatocyte nuclear factor 3 and liver factor B1 are differently affected by pure and partial antiestrogens. *Eur. J. Biochem.* 225, 441-448.
- Drewes,T., Senkel,S., Holewa,B., and Ryffel,G.U. (1996). Human hepatocyte nuclear factor 4 isoforms are encoded by distinct and differentially expressed genes. *Mol. Cell Biol.* 16, 925-931.
- Duncan,S.A., Nagy,A., and Chan,W. (1997). Murine gastrulation requires HNF4 regulated gene expression in the visceral endoderm: tetraploid rescue of HNF4(-/-) embryos. *Development* 124, 279-287.
- Duncan,S.A., Navas,M.A., Dufort,D., Rossant,J., and Stoffel,M. (1998). Regulation of a transcription factor network required for differentiation and metabolism. *Science* 281, 692-695.
- Ellard,S. (2000). Hepatocyte nuclear factor 1 alpha (HNF1 $\alpha$ ) mutations in maturity onset diabetes of the young. *Hum. Mutat.* 16, 377-385.
- Emens,L.A., Landers,D.W., and Moss,L.G. (1992). Hepatocyte nuclear factor 1 alpha is expressed in a hamster insulinoma line and transactivates the rat insulin I gene. *Proc. Natl. Acad. Sci. U. S. A.* 89, 7300-7304.
- Evan,G.I., Lewis,G.K., Ramsay,G., and Bishop,J.M. (1985). Isolation of monoclonal antibodies specific for human c-myc proto-oncogene product. *Mol. Cell Biol.* 5, 3610-3616.
- Fearon,E.R. (1997). Human cancer syndromes: Clues to the origin and nature of cancer. *Science* 278, 1043-1050.

- Finegood,D.T., Scaglia,L., and Bonner-Weir,S. (1995). Dynamics of beta-cell mass in the growing rat pancreas. Estimation with a simple mathematical model. *Diabetes* 44, 249-256.
- Froguel,P. and Velho,G. (1999). Molecular genetics of maturity onset diabetes of the young. *Trends Endocrinol. Metab* 10, 142-146.
- Fuhrmann-Benzakein,E., Garcia-Gabay,I., Pepper,M.S., Vassalli,J.D., and Herrera,P.L. (2000). Inducible and irreversible control of gene expression using a single transgene. *Nucleic Acids Res.* 28, E99.
- Furuta,H., Iwasaki,N., Oda,N., Hinokio,Y., Horikawa,Y., Yamagata,K., Yano,N., Sugahiro,J., Ogata,M., Ohgawara,H., Omori,Y., Iwamoto,Y., and Bell,G.I. (1997). Organization and partial sequence of the hepatocyte nuclear factor 4 alpha/MODY1 gene and identification of a missense mutation, R127W, in a Japanese family with MODY. *Diabetes* 46, 1652-1657.
- Glass,C.K. and Rosenfeld,M.G. (2000). The coregulator exchange in transcriptional functions of nuclear receptors. *Genes Dev.* 14, 121-141.
- Goodman,R.H. and Smolik,S. (2000). CBP/p300 in cell growth, transformation, and development. *Genes Dev.* 14, 1553-1577.
- Gragnoli,C., Lindner,T., Cockburn,B.N., Kaisaki,P.J., Gragnoli,F., Marozzi,G., and Bell,G.I. (1997). Maturity onset diabetes of the young due to a mutation in the hepatocyte nuclear factor 4 alpha binding site in the promoter of the hepatocyte nuclear factor 1 alpha gene. *Diabetes* 46, 1648-1651.
- Hadzopoulou-Cladaras,M., Kistanova,E., Evagelopoulou,C., Zeng,S.Y., Cladaras,C., and Ladias,J.A.A. (1997). Functional domains of the nuclear receptor hepatocyte nuclear factor 4. *J. Biol. Chem.* 272, 539-550.
- Hani,E.H., Suaud,L., Boutin,P., Chevre,J.C., Durand,E., Philippi,A., Demenais,F., Vionnet,N., Furuta,H., Velho,G., Bell,G.I., Laine,B., and Froguel,P. (1998). A missense mutation in hepatocyte nuclear factor 4 alpha, resulting in a reduced transactivation activity, in human late onset non insulin dependent diabetes mellitus. *J. Clin. Invest.* 101, 521-526.
- Hattersley,A.T. (1998). Maturity onset diabetes of the young: clinical heterogeneity explained by genetic heterogeneity. *Diabet. Med.* 15, 15-24.
- Hertz,R., Magenheimer,J., Berman,I., and Bar-Tana,J. (1998). Fatty acyl-CoA thioesters are ligands of hepatic nuclear factor 4 alpha. *Nature* 392, 512-516.
- Holewa,B., Pogge v.Strandmann,E., Zapp,D., Lorenz,P., and Ryffel,G.U. (1996). Transcriptional hierarchy in *Xenopus* embryogenesis: HNF4 a maternal factor involved in the developmental activation of the gene encoding the tissue specific transcription factor HNF1 $\alpha$  (LFB1). *Mech. Dev.* 54, 45-57.
- Holewa,B., Zapp,D., Drewes,T., Senkel,S., and Ryffel,G.U. (1997). HNF4 $\beta$ , a new gene of the HNF4 family with distinct activation and expression profiles in oogenesis and embryogenesis of *Xenopus laevis*. *Mol. Cell Biol.* 17, 687-694.
- Ito,M., Achermann,J.C., and Jameson,J.L. (2000). A naturally occurring steroidogenic factor 1 mutation exhibits differential binding and activation of target genes. *J. Biol. Chem.* 275, 31708-31714.

- Jiang,G., Nepomuceno,L., Hopkins,K., and Sladek,F.M. (1995). Exclusive homodimerization of the orphan receptor hepatocyte nuclear factor 4 defines a new subclass of nuclear receptors. *Mol. Cell Biol.* *15*, 5131-5143.
- Jiang,G. and Sladek,F.M. (1997). The DNA binding domain of hepatocyte nuclear factor 4 mediates cooperative, specific binding to DNA and heterodimerization with the retinoid X receptor alpha. *J. Biol. Chem.* *272*, 1218-1225.
- Kapitskaya,M.Z., Dittmer,N.T., Deitsch,K.W., Cho,W.L., Taylor,D.G., Leff,T., and Raikhel,A.S. (1998). Three isoforms of a hepatocyte nuclear factor 4 transcription factor with tissue and stage specific expression in the adult mosquito. *J. Biol. Chem.* *273*, 29801-29810.
- Knowles,B.B., Howe,C.C., and Aden,D.P. (1980). Human hepatocellular carcinoma cell lines secrete the major plasma proteins and hepatitis B surface antigen. *Science* *209*, 497-499.
- Ktistaki,E. and Talianidis,I. (1997). Chicken ovalbumin upstream promoter transcription factors act as auxiliary cofactors for hepatocyte nuclear factor 4 and enhance hepatic gene expression. *Mol. Cell Biol.* *17*, 2790-2797.
- Kumar,R. and Thompson,E.B. (1999). The structure of the nuclear hormone receptors. *Steroids* *64*, 310-319.
- Kung,A.L., Rebel,V.I., Bronson,R.T., Ch'ng,L.E., Sieff,C.A., Livingston,D.M., and Yao,T.P. (2000). Gene dose dependent control of hematopoiesis and hematologic tumor suppression by CBP. *Genes Dev.* *14*, 272-277.
- Kuo,C.J., Conley,P.B., Chen,L., Sladek,F.M., Darnell,J.E., Jr., and Crabtree,G.R. (1992). A transcriptional hierarchy involved in mammalian cell type specification. *Nature* *355*, 457-461.
- Laine,B., Eeckhoute,J., Suaud,L., Briche,I., Furuta,H., Bell,G.I., and Formstecher,P. (2000). Functional properties of the R154X HNF4 $\alpha$  protein generated by a mutation associated with maturity onset diabetes of the young, type 1. *FEBS Lett.* *479*, 41-45.
- Laudet,V., Hänni,C., Coll,J., Catzefis,F., and Stehelin,D. (1992). Evolution of the nuclear receptor gene superfamily. *EMBO J.* *11*, 1003-1013.
- Lausen,J., Thomas,H., Lemm,I., Bulman,M., Borgschulze,M., Lingott,A., Hattersley,A.T., and Ryffel,G.U. (2000). Naturally occurring mutations in the human HNF4 $\alpha$  gene impair the function of the transcription factor to a varying degree. *Nucleic Acids Res.* *28*, 430-437.
- Lee,Y.H., Sauer,B., and Gonzalez,F.J. (1998). Laron dwarfism and non insulin dependent diabetes mellitus in the HNF1 $\alpha$  knockout mouse. *Mol. Cell Biol.* *18*, 3059-3068.
- Lehto,M., Bitzen,P.O., Isomaa,B., Wipemo,C., Wessman,Y., Forsblom,C., Tuomi,T., Taskinen,M.R., and Groop,L. (1999a). Mutation in the HNF4 $\alpha$  gene affects insulin secretion and triglyceride metabolism. *Diabetes* *48*, 423-425.
- Lehto,M., Wipemo,C., Ivarsson,S.A., Lindgren,C., Lipsanen-Nyman,M., Weng,J., Wibell,L., Widen,E., Tuomi,T., and Groop,L. (1999b). High frequency of mutations in MODY and mitochondrial genes in Scandinavian patients with familial early onset diabetes. *Diabetologia* *42*, 1131-1137.

- Li, J., Ning, G., and Duncan, S.A. (2000). Mammalian hepatocyte differentiation requires the transcription factor HNF4 $\alpha$ . *Genes Dev.* *14*, 464-474.
- Lindblad, P., Chow, W.H., Chan, J., Bergstrom, A., Wolk, A., Gridley, G., McLaughlin, J.K., Nyren, O., and Adami, H.O. (1999). The role of diabetes mellitus in the aetiology of renal cell cancer. *Diabetologia* *42*, 107-112.
- Lindner, T., Gragnoli, C., Furuta, H., Cockburn, B.N., Petzold, C., Rietzsch, H., Weiss, U., Schulze, J., and Bell, G.I. (1997). Hepatic function in a family with a nonsense mutation (R154X) in the hepatocyte nuclear factor 4 alpha/MODY1 gene. *J. Clin. Invest* *100*, 1400-1405.
- Love, J.D., Gooch, J.T., Nagy, L., Chatterjee, K.K., and Schwabe, J.W.R. (2000). Transcriptional repression by nuclear receptors: Mechanism and role in disease. *Biochemical Society Transactions* *28*, 390-396.
- Mader, S., Kumar, V., de Verneuil, H., and Chambon, P. (1989). Three amino acids of the oestrogen receptor are essential to its ability to distinguish an oestrogen from a glucocorticoid responsive element. *Nature* *338*, 271-274.
- Malecki, M.T., Yang, Y., Antonellis, A., Curtis, S., Warram, J.H., and Krolewski, A.S. (1999). Identification of new mutations in the hepatocyte nuclear factor 4 alpha gene among families with early onset Type 2 diabetes mellitus. *Diabet. Med.* *16*, 193-200.
- Malloy, P.J., Pike, J.W., and Feldman, D. (1999). The vitamin D receptor and the syndrome of hereditary 1,25-dihydroxyvitamin-D resistant rickets. *Endocr. Rev.* *20*, 156-188.
- Mangelsdorf, D.J. and Evans, R.M. (1995). The RXR heterodimers and orphan receptors. *Cell* *83*, 841-850.
- Mangelsdorf, D.J., Thummel, C., Beato, M., Herrlich, P., Schutz, G., Umesono, K., Blumberg, B., Kastner, P., Mark, M., and Chambon, P. (1995). The nuclear receptor superfamily: the second decade. *Cell* *83*, 835-839.
- Massague, J., Blain, S.W., and Lo, R.S. (2000). TGF $\beta$  signaling in growth control, cancer and heritable disorders. *Cell* *103*, 295-309.
- McKenna, N.J., Lanz, R.B., and O'Malley, B.W. (1999). Nuclear receptor coregulators: cellular and molecular biology. *Endocr. Rev.* *20*, 321-344.
- McNair, A., Cereghini, S., Brand, H., Smith, T., Breillat, C., and Gannon, F. (2000). Synergistic activation of the atlantic salmon hepatocyte nuclear factor (HNF) 1 promoter by the orphan nuclear receptors HNF4 and chicken ovalbumin upstream promoter transcription factor I (COUP-TFI). *Biochem. J.* *352 Pt 2*, 557-564.
- Miquerol, L., Lopez, S., Cartier, N., Tulliez, M., Raymondjean, M., and Kahn, A. (1994). Expression of the L-type pyruvate kinase gene and the hepatocyte nuclear factor 4 transcription factor in exocrine and endocrine pancreas. *J. Biol. Chem.* *269*, 8944-8951.
- Möller, A.M., Dalgaard, L.T., Ambye, L., Hansen, L., Schmitz, O., Hansen, T., and Pedersen, O. (1999). A novel Phe75fsdelT mutation in the hepatocyte nuclear factor 4 alpha gene in a Danish pedigree with maturity onset diabetes of the young. *J. Clin. Endocrinol. Metab.* *84*, 367-369.

- Möller,A.M., Urhammer,S.A., Dalgaard,L.T., Reneland,R., Berglund,L., Hansen,T., Clausen,J.O., Lithell,H., and Pedersen,O. (1997). Studies of the genetic variability of the coding region of the hepatocyte nuclear factor 4 alpha in caucasians with maturity onset NIDDM. *Diabetologia* 40, 980-983.
- Morrisey,E.E., Tang,Z., Sigrist,K., Lu,M.M., Jiang,F., Ip,H.S., and Parmacek,M.S. (1998). GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev.* 12, 3579-3590.
- Nakhei,H., Lingott,A., Lemm,I., and Ryffel,G.U. (1998). An alternative splice variant of the tissue specific transcription factor HNF4 $\alpha$  predominates in undifferentiated murine cell types. *Nucleic. Acids. Res.* 26, 497-504.
- Nastos,A., Pogge v.Strandmann,E., Weber,H., and Ryffel,G.U. (1998). The embryonic expression of the tissue specific transcription factor HNF1 $\alpha$  in *Xenopus*: rapid activation by HNF4 and delayed induction by mesoderm inducers. *Nucleic. Acids. Res.* 26, 5602-5608.
- Navas,M.A., Munoz-Elias,E.J., Kim,J., Shih,D., and Stoffel,M. (1999). Functional characterization of the MODY1 gene mutations HNF4(R127W), HNF4(V255M), and HNF4(E276Q). *Diabetes* 48, 1459-1465.
- Nuclear Receptors Nomenclature Committee 1999 (1999). A unified nomenclature system for the nuclear receptor superfamily. *Cell* 97, 161-163.
- Okita,K., Yang,Q., Yamagata,K., Hangenfeldt,K.A., Miyagawa,J., Kajimoto,Y., Nakajima,H., Namba,M., Wollheim,C.B., Hanafusa,T., and Matsuzawa,Y. (1999). Human insulin gene is a target gene of hepatocyte nuclear factor 1 alpha (HNF1 $\alpha$ ) and HNF1 $\beta$ . *Biochem. Biophys. Res. Commun.* 263, 566-569.
- Peiler,G., Bockmann,B., Nakhei,H., and Ryffel,G.U. (2000). Inhibitor of the tissue-specific transcription factor HNF4, a potential regulator in early *Xenopus* development. *Mol. Cell Biol.* 20, 8676-8683.
- Plengvidhya,N., Antonellis,A., Wogan,L.T., Poleev,A., Borgschulze,M., Warram,J.H., Ryffel,G.U., Krolewski,A.S., and Doria,A. (1999). Hepatocyte nuclear factor 4 gamma: cDNA sequence, gene organization, and mutation screening in early onset autosomal-dominant type 2 diabetes. *Diabetes* 48, 2099-2102.
- Price,J.A., Fossey,S.C., Sale,M.M., Brewer,C.S., Freedman,B.I., Wuerth,J.P., and Bowden,D.W. (2000). Analysis of the HNF4 $\alpha$  gene in Caucasian type-II diabetic nephropathic patients. *Diabetologia* 43, 364-372.
- Reijnen,M.J., Sladek,F.M., Bertina,R.M., and Reitsma,P.H. (1992). Disruption of a binding site for hepatocyte nuclear factor 4 results in hemophilia B Leyden. *Proc. Natl. Acad. Sci. U. S. A.* 89, 6300-6303.
- Rempel,A., Mathupala,S.P., Griffin,C.A., Hawkins,A.L., and Pedersen,P.L. (1996). Glucose catabolism in cancer cells: amplification of the gene encoding type-II hexokinase. *Cancer Res.* 56, 2468-2471.

- Rosen,E.D., Walkey,C.J., Puigserver,P., and Spiegelman,B.M. (2000). Transcriptional regulation of adipogenesis. *Genes Dev.* *14*, 1293-1307.
- Rupp,R.A., Snider,L., and Weintraub,H. (1994). *Xenopus* embryos regulate the nuclear localization of XMyoD. *Genes Dev.* *8*, 1311-1323.
- Ryffel,G.U. and Lingott,A. (2000). Distinct promoter elements mediate endodermal and mesodermal expression of the HNF1 $\alpha$  promoter in transgenic *Xenopus*. *Mech. Dev.* *90*, 65-75.
- Sambrock,J., Fritsch,E.F., and Maniatis,T. (1989). Molecular cloning - a laboratory manual. (2<sup>nd</sup> edition) Cold Spring Harbour Laboratory, Cold Spring Harbour, New York.
- Sarraf,P., Mueller,E., Smith,W.M., Wright,H.M., Kum,J.B., Aaltonen,L.A., de la,C.A., Spiegelman,B.M., and Eng,C. (1999). Loss of function mutations in PPAR $\gamma$  associated with human colon cancer. *Mol. Cell* *3*, 799-804.
- Sel,S., Ebert,T., Ryffel,G.U., and Drewes,T. (1996). Human renal cell carcinogenesis is accompanied by a coordinate loss of the tissue specific transcription factors HNF4 $\alpha$  and HNF1 $\alpha$ . *Cancer Lett.* *101*, 205-210.
- Shih,D.Q., Dansky,H.M., Fleisher,M., Assmann,G., Fajans,S.S., and Stoffel,M. (2000). Genotype/phenotype relationships in HNF4 $\alpha$ /MODY1: haploinsufficiency is associated with reduced apolipoprotein (AII), apolipoprotein (CIII), lipoprotein(a), and triglyceride levels. *Diabetes* *49*, 832-837.
- Sladek,F.M., Dallas-Yang,Q., and Nepomuceno,L. (1998). MODY1 mutation Q268X in hepatocyte nuclear factor 4 alpha allows for dimerization in solution but causes abnormal subcellular localization. *Diabetes* *47*, 985-990.
- Sladek,F.M. and Seidel,S.D. (2001). Hepatocyte nuclear factor 4 alpha. In: Nuclear receptors and genetic disease, T.P.Burris and E.R.B.McCabe (eds.), Academic Press, pp. 309-348.
- Sladek,F.M., Zhong,W.M., Lai,E., and Darnell,J.E., Jr. (1990). Liver-enriched transcription factor HNF4 is a novel member of the steroid hormone receptor superfamily. *Genes Dev.* *4*, 2353-2365.
- Spaeth,G.F. and Weiss,M.C. (1997). Hepatocyte nuclear factor 4 expression overcomes repression of the hepatic phenotype in dedifferentiated hepatoma cells. *Mol. Cell Biol.* *17*, 1913-1922.
- Stoffers,D.A., Thomas,M.K., and Habener,J.F. (1997a). Homeodomain protein IDX-1 a master regulator of pancreas development and insulin gene expression. *Trends. Endocrinol. Metab.* *8*, 145-151.
- Stoffers,D.A., Ferrer,J., Clarke,W.L., and Habener,J.F. (1997b). Early onset type-II diabetes mellitus (MODY4) linked to IPF1. *Nat. Genet.* *17*, 138-139.
- Stoffers,D.A., Zinkin,N.T., Stanojevic,V., Clarke,W.L., and Habener,J.F. (1997c). Pancreatic agenesis attributable to a single nucleotide deletion in the human IPF1 gene coding sequence. *Nat. Genet.* *15*, 106-110.

- Suaud,L., Formstecher,P., and Laine,B. (1999a). The activity of the activation function 2 of the human hepatocyte nuclear factor 4 (HNF4 $\alpha$ ) is differently modulated by F domains from various origins. *Biochem. J.* 340, 161-169.
- Suaud,L., Hemimou,Y., Formstecher,P., and Laine,B. (1999b). Functional study of the E276Q mutant hepatocyte nuclear factor 4 alpha found in type 1 maturity onset diabetes of the young: impaired synergy with chicken ovalbumin upstream promoter transcription factor II on the hepatocyte nuclear factor 1 promoter. *Diabetes* 48, 1162-1167.
- Swevers,L. and Iatrou,K. (1998). The orphan receptor BmHNF4 of the silkworm *Bombyx mori*: ovarian and zygotic expression of two mRNA isoforms encoding polypeptides with different activating domains. *Mech. Dev.* 72, 3-13.
- Torchia,J., Glass,C., and Rosenfeld,M.G. (1998). Co-activators and co-repressors in the integration of transcriptional responses. *Curr. Opin. Cell Biol.* 10, 373-383.
- Tronche,F., Ringeisen,F., Blumenfeld,M., Yaniv,M., and Pontoglio,M. (1997). Analysis of the distribution of binding sites for a tissue specific transcription factor in the vertebrate genome. *J. Mol. Biol.* 266, 231-245.
- Tronche,F. and Yaniv,M. (1992). HNF1, a homeoprotein member of the hepatic transcription regulatory network. *Bioessays* 14, 579-587.
- Umesono,K. and Evans,R.M. (1989). Determinants of target gene specificity for steroid/thyroid hormone receptors. *Cell* 57, 1139-1146.
- Vaxillaire,M., Abderrahmani,A., Boutin,P., Bailleul,B., Froguel,P., Yaniv,M., and Pontoglio,M. (1999). Anatomy of a homeoprotein revealed by the analysis of human MODY3 mutations. *J. Biol. Chem.* 274, 35639-35646.
- Wang,H., Maechler,P., Antinozzi,P.A., Hagenfeldt,K.A., and Wollheim,C.B. (2000a). HNF4 $\alpha$  regulates the expression of pancreatic beta-cell genes implicated in glucose metabolism and nutrient induced insulin secretion. *J. Biol. Chem.* 275, 35953-35959.
- Wang,H., Antinozzi,P.A., Hagenfeldt,K.A., Maechler,P., and Wollheim,C.B. (2000b). Molecular targets of a human HNF1 $\alpha$  mutation responsible for pancreatic beta-cell dysfunction. *EMBO J.* 19, 4257-4264.
- Wang,H., Maechler,P., Hagenfeldt,K.A., and Wollheim,C.B. (1998). Dominant-negative suppression of HNF1 $\alpha$  function results in defective insulin gene transcription and impaired metabolism secretion coupling in a pancreatic beta-cell line. *EMBO J.* 17, 6701-6713.
- Weber,H., Holewa,B., Jones,E.A., and Ryffel,G.U. (1996). Mesoderm and endoderm differentiation in animal cap explants: identification of the HNF4 binding site as an activin A responsive element in the *Xenopus* HNF1 $\alpha$  promoter. *Development* 122, 1975-1984.
- Wild,W., Pogge,v.S., Nastos,A., Senkel,S., Lingott-Frieg,A., Bulman,M., Bingham,C., Ellard,S., Hattersley,A.T., and Ryffel,G.U. (2000). The mutated human gene encoding hepatocyte nuclear factor 1 beta inhibits kidney formation in developing *Xenopus* embryos. *Proc. Natl. Acad. Sci. U. S. A* 97, 4695-4700.
- Winter,W.E. and Silverstein,J.H. (2000). Molecular and genetic bases for maturity onset diabetes of youth. *Curr. Opin. Pediatr.* 12, 388-393.



- Yamagata,K., Furuta,H., Oda,N., Kaisaki,P.J., Menzel,S., Cox,N.J., Fajans,S.S., Signorini,S., Stoffel,M., and Bell,G.I. (1996). Mutations in the hepatocyte nuclear factor 4 alpha gene in maturity onset diabetes of the young (MODY1). *Nature* 384, 458-460.
- Yamaoka,T. and Itakura,M. (1999). Development of pancreatic islets. *Int. J. Mol. Med.* 3, 247-261.
- Yang,Q., Yamagata,K., Yamamoto,K., Cao,Y., Miyagawa,J., Fukamizu,A., Hanafusa,T., and Matsuzawa,Y. (2000). R127W-HNF4 $\alpha$  is a loss of function mutation but not a rare polymorphism and causes Type-II diabetes in a Japanese family with MODY1. *Diabetologia* 43, 520-524.
- Zapp,D., Bartkowski,S., Holewa,B., Zoidl,C., Klein-Hitpass,L., and Ryffel,G.U. (1993). Elements and factors involved in tissue specific and embryonic expression of the liver transcription factor LFB1 in *Xenopus laevis*. *Mol. Cell Biol.* 13, 6416-6426.
- Zhong,W., Mirkovitch,J., and Darnell,J.E., Jr. (1994). Tissue specific regulation of mouse hepatocyte nuclear factor 4 expression. *Mol. Cell Biol.* 14, 7276-7284.