Frontal and temporal sources of mismatch negativity in healthy controls, patients at onset of schizophrenia in adolescence and others at 15 years after onset.

Oknina, L. B.^{a,1}, Wild-Wall, N.^{b,1}, Oades, R. D.^b, Juran, S. A.^b, Röpcke, B.^b, Pfueller, U.^c, Weisbrod, M.^c, Chan, E.^d, Chen, E. Y. H.^d

2005 Schizophrenia Research, 76, 25-41.

Biopsychology Group, University Clinic for Child and Adolescent Psychiatry, Virchowstr. 174, 45147, Essen, Germany.

- a) Institute of Higher Nervous Activity & Neurophysiology, Burdenko Neurosurgery Institute, Butlerova Str. 5a, Moscow, Russia.
- b) University Clinic for Child and Adolescent Psychiatry, Virchowstr. 174, 45147, Essen, Germany.
- c) Section Experimental Psychopathology, University Psychiatry Clinic, Voßstr 6, 69115 Heidelberg, Germany.
- d) Queen Mary Hospital, University of Hong Kong, Pokfulam Road, Hong Kong, PRC.
- 1) Both authors contributed equally to this work

Abstract

Mismatch negativity (MMN) is an event-related potential measure of auditory change detection. It is widely reported to be smaller in patients with schizophrenia and may not improve along with otherwise successful clinical treatment. The main aim of this report is to explore ways of measuring and presenting four features of frequency-deviant MMN dipole sources (dipole moment, peak latency, brain location and orientation) and to relate these to the processes of psychopathology and illness progression. Data from early onset patients (EOS) at the start of the illness in adolescence, and others who had their first break in adolescence 15 years ago (S-15Y) were compared with two groups of age-matched healthy controls (C-EOS, C-15Y). A 4-source model fitted the MMN waveform recorded from all four groups, whether MMN amplitude was more (EOS) or less (S-15Y) reduced. The locations were in the left superior temporal and anterior cingulate gyri, right superior temporal and inferior/mid frontal cortices. Dipole latencies confirmed a bottom-up sequence of processing and dipole moments were larger in the temporal lobes and on the left. Patients showed small dipole location changes that were more marked in the S-15Y than the EOS group (more rostral for the left anterior cingulate, more caudal for the right mid-frontal dipole) consistent with illness progression. The modelling of MMN dipole sources on brain atlas and anatomical images suggests that there is a degree of dissociation during illness between small progressive anatomical changes and some functional recovery indexed by scalp recordings from patients with an onset in adolescence 15 years before compared to adolescents in their first episode.

Key Words:

Attention, Cingulate, Dipole, Frontal cortex, Mismatch Negativity (MMN), Schizophrenia

Correspondence:

Robert D. Oades, University Clinic for Child and Adolescent Psychiatry, Virchowstr. 174, 45147 Essen, Germany.

Acknowledgments:

LBO and NWW were supported by the Alfried Krupp von Bohlen und Halbach Stiftung. We gratefully acknowledge the help of V. Cheung, K. Chum, O. C. Chan, K. Herwig, R. Torres, J. Sachsse, R. Franzka and R. Windelschmitt.

1. Introduction

The ability to detect a change of stimulation from the steady ongoing situation is crucial to

being able to assign value to the change and organise a potentially appropriate response to it. Changes to a sequence of auditory stimuli can be

registered automatically in the brain and recorded as an event-related potential (ERP) known as the mismatch negativity (MMN: Näätänen et al., 1978).

Dipole sources for MMN have been reported in the temporal cortices bilaterally, the right inferior frontal and the left anterior cingulate cortex (Jemel et al., 2002; Valkonen-Korhonen et al., 2003). The auditory cortical sources may represent the establishment of a sensory memory trace for the status quo (Paavilainen et al., 1991; Jacobsen and Schröger, 2001) while the inferior/mid-frontal source may be part of a region involved in generic change detection (Beck et al., 2001). The latter is seen as a pre-requisite to switching from the ongoing automatic mode of information processing and bringing top-down controlled attentional processing into effect (Schröger 1996; Escera et al., 1998). An alternative but not exclusive interpretation could invoke an orienting mechanism to explain the right frontal advantage in auditory cued attention in ERP work (Ofek and Pratt, 2004) and f-MRI studies of visual attention (Arrington et al., 2000). The activity of the preceding left cingulate source may reflect the registration of the contrast between similar and different stimuli necessary to elicit the right frontal “switch”. Such a role may be analogous to the reported role of the cingulate in conflict monitoring and error detection (Carter et al., 1998; Brazdil et al., 2002). Both frontal roles are fundamental to numerous cognitive functions reported to be impaired in schizophrenia.

The MMN mechanism (or a part thereof) is widely acknowledged to be impaired specifically in patients with schizophrenia (Oades 1995; Javitt et al., 1995; Umbricht et al., 2003). Studies using magnetoencephalography (MEG) report reduced temp-oral lobe dipole strength, especially in the left hemisphere (Kircher et al., 2004; Kasai et al., 2003; Pekkonen et al., 2002). But there is only one similar report based on ERPs (Youn et al., 2003). MEG is insensitive to frontal sources, but indirect evidence from ERP recordings has indicated an impaired frontal contribution to MMN, specific to schizophrenia (i.e., a reduced positive component recorded from mastoid sites, Todd et al., 2002; Sato et al., 2003). Some laboratories report that the MMN impairment is more marked for duration deviants (Michie et al., 2000; Umbricht et al., 2003), others find a more

marked effect with variations of stimulus frequency (Shinozaki et al., 2002; Oades et al., 1996). This report focuses on the latter and the putative contribution from frontal sources.

There is controversy over whether anatomical or functional impairments progress with the course of illness. Some authors find that impairments develop with illness duration (Salisbury et al., 1999; Umbricht et al., 2002). But others report deficits at onset in adults (Shinozaki et al., 2002) and in adolescents with recent onset of schizophrenia, that continue into the later phases of the illness (Oades et al., 1997). The current study investigated both young patients at onset and a second group about 15 years after illness-onset in adolescence to try to resolve some of these structural and functional issues. First we calculated the potential location of MMN dipole sources in two groups of patients with schizophrenia on brain atlas and high-resolution MR-images of the frontal/temporal lobes. Secondly we sought to confirm that predicted changes in frontal lobe sources also occur using frequency deviant stimuli. Thirdly we investigated whether such differences could be demonstrated in patients experiencing their first break or whether reductions of MMN improve or deteriorate in patients some 15 years after their first episode.

2. Methods

2.1 Subjects

At the time-point taken in the ongoing longitudinal study for this cross-sectional analysis there were 71 subjects: 25 young patients hospitalised with a first episode of schizophrenia (mean age 17.6 years, SD 2.3), 17 patients who had had their first admission for schizophrenia in the Essen Clinic about 15 years before, as adolescents (mean 14.4 years, SD 3.5 range 8.1-19.7 years: mean age 32.3 years SD 3.6). There were 14 healthy adolescents (mean age 17.2 years, SD 1.8), and 15 healthy adults in the comparison groups (mean age 30.0 years, SD 6.0).

Age-matched controls were recruited by advertisement, and paid for their participation. They were matched to the patients of their age group on the socio-economic scale (parent profession) to within one point on a scale of 6 and, like the patients, they were all right- or mixed handed (Table 1). Based on the results of a semi-structured interview, none of the

adolescent controls (C-EOS) for the early-onset (C-15Y) for the group diagnosed with schizophrenia group (EOS) nor the adult controls

Table 1 **Characteristics of Subject Groups (mean + SD)**

	1 st -episode Inpatients EOS		Adolescent Controls C-EOS		Adult Outpatients S-15Y		Adult Controls C-15Y	
	EOS	SD	C-EOS	SD	S-15Y	SD	C-15Y	SD
Subjects, Male/Female	19 / 6		8 / 6		11 / 6		6 / 9	
Age (years)	17.6	2.4	17.2	1.8	32.3	3.6	30.0	6.1
Handedness ¹	93.5	9.0	70.8	42.3	75.0	50.1	79.2	26.7
SES (parent occupation ²)	3.6	1.4	2.8	0.6	4.2	1.4	3.3	1.4
Short-IQ ³	95	20.7	113	7.3	91	17.3	115	17.5
Diagnosis (at test: DSMIV) --								
Paranoid	19				8			
Disorganised	4				1			
Undifferentiated	1				1			
Residual	0				5			
Schizoaffective	0				2			
Antipsychotic Medication ³	609	335	-	-	366	127		
Without medication (N)	11	-	-	-	3	-	-	-
N1 standard latency (ms)	98	6	101	10	96	7	97	11
N1 deviant latency (ms)	113	18	108	11	112	20	103	11
MMN trials accepted	64	8	62	10	64	13	64	10
MMN latency (ms)	137	27	140	22	135	31	135	34
MMN amplitude (μV) --								
FCz (windows 1/2/3)	-1.35/-1.07*/-0.49		-1.42/-1.51/-0.43		-1.68/-1.12 /-0.44		-1.58/-0.97/-0.87	
M1 (windows 1/2/3)	1.39/ 1.19 ⁺ / 0.55		1.77/ 1.76/ 1.25		1.19/ 0.19* / -0.25 [#]		1.55/ 1.44/ 1.03	
M2 (windows 1/2/3)	1.65/ 1.06 [#] / 0.35*		2.16/ 2.18/ 1.34		1.42/ 0.50 / -0.06*		1.62/ 0.94/ 0.65	

¹Edinburgh handedness inventory (-100 [left-handed] to +100 [right-handed] Oldfield, 1971). ²The socio-economic scale (SES, ratings 1-6, Erikson and Goldthorpe 1992). Group differences were not significant. ³Short-IQ based on 4 tests (information, arithmetic, digit-symbol and block-design): patient scores were lower than the matched controls (t -2.7 to -3.7, $p < 0.02$). ³ CPZ mg/day: Chlorpromazine equivalents for those on medication (see text). There were no significant group differences for the ERP latencies at FCz: for N1 amplitudes see text. MMN amplitudes are shown for 3 frontal and mastoid electrode sites in three 30 ms time windows (1, 105-135; 2, 135-165; 3 165-195 ms: SD range 0.24-0.46). ⁺ $p=0.09$, * $p=0.02-0.04$, [#] $p=0.006-0.01$

schizophrenia 15y ago (S-15Y) had been ill or had a history of neurological or psychiatric illness or used drugs affecting the central nervous system. They were screened using the SCID-II and excluded if there were lifetime or current DSM-IV Axis II diagnoses, including schizotypy.

EOS patients were admitted consecutive-ly for functional psychosis. The initial diagnosis was made by the senior ward physician. They were re-examined for entry to the study and the diagnosis confirmed after at least six months to exclude affective, schizoaffective and schizophreniform psychoses. Patients were admitted to the S-15Y group on the basis of their initial diagnosis of

schizophrenia 15 years before: (a current diagnosis of schizoaffective disorder was accepted in 2 cases). Patients were screened to exclude other major psychiatric or somatic illness, alcohol abuse in the last 5 years and current substance abuse other than nicotine. Schizophrenia subtypes were defined by DSM-IV criteria, whereby the undifferentiated type was regarded as a residual category that contrasts with the paranoid, disorganized and catatonic subtypes (see Table 1).

S-15Y patients were recruited from 34 in outpatient care (see Eggers et al. (1999). Eleven (65%) had a partial remission (CGI 3-5) and 6 (35%)

a chronic course or no remission (CGI 6-8). The main symptoms were emotional and social withdrawal, reduced affect and increased anxiety and tension: the dominant positive symptom was 'excitement'. Seven (41%) had severe impairments in several areas of social function (GAS < 51) and 11 (59%) had moderate impairments in at least one (GAS 51-70). Since onset they had had a mean of 4.6 hospitalizations (range 1-12). They did not differ from the whole group on age, gender, severity of illness or social function. Reasons for non-participation included severe psychopathology, but 4 were considered as fully remitted.

Following agreement of the University-Clinics' ethics committee to the protocol (following the Declaration of Helsinki), the study was undertaken with the understanding of the subjects and their care-givers who both provided written consent to participate.

2.2 Medication

EOS patients were tested in the first 2 weeks of their stay, contingent on the responsible physician's opinion that they were capable of performing the tests, and medication had been stable for at least 3 days. Antipsychotic drugs were administered according to the clinical requirements. Doses were converted to chlorpromazine equivalents (CPZ) according to Benkert and Hippus (1986) Rey et al. (1989), Kane (1996) and information from the suppliers of olanzapine and quetiapine. Eleven EOS patients were without medication at testing, two were given typical antipsychotic drugs (haloperidol, fluphenazine), 12 received atypical medication (clozapine, olanzapine, risperidone, amisulpride or quetiapine: CPZ 100-1300). In the S-15Y group 3 patients were without medication, two received typical and 12 received atypical antipsychotic drugs (CPZ 200-720: Table 1).

2.3 Stimulus Presentation, EEG Recording and MR-Imaging

All subjects had normal hearing (BCA audiometer (low-frequency thresholds < 30dB, inter-aural differences <10dB) and had normal or corrected-to-normal vision. A binaural 3-tone auditory oddball presentation consisted of sinusoidal tones (76 dBspl). The standards had a frequency of 800 Hz ($p = 80\%$, duration 80 ms, 10 ms rise/fall times), deviants had either a frequency of 600 Hz ($p = 10\%$) or a duration of 40

ms ($p=10\%$). The stimulus onset asynchrony (SOA) varied from 850-1050 ms (mean 950 ms). Salient deviants were chosen to produce large amplitudes with more reliable test-retest sensitivity and signal-to-noise ratios for localisation. A relatively long SOA was used to facilitate response in the subsequent auditory discrimination, (data not presented.) The order of stimulus presentation was pseudo-random with each deviant preceded by at least one standard. The MMN was calculated from the frequency-deviant ERP minus standard ERP waveforms.

Four auditory trial blocks (400 tones/ block) were run during a simple visual red or green circle discrimination (50:50). The circles subtended an angle of 3.8° at 1.5 m and alternated randomly every 1100 ms on a PC screen. Subjects responded alternately between blocks with the left and right hand to the green circle. Auditory stimulus detection is not suppressed during various concurrent visual processes (Harris and Lieberman, 1996), but the visual and auditory stimulus onsets were controlled so that they did not coincide.

The EEG was recorded (Neuroscan) from 32 electrodes in an extended 10-20 system (Electrocap). These were the Fpz, Fz, Cz, Pz, Oz, F3, F7, F4, F8, C3, C4, P3, P4, T3, T4, T5, T6, both mastoids, FCz, CPz (longitudinal axis), FT7-FT8 (fronto-temporal), FC3-FC4 (fronto-central), CP3-CP4 (centro-parietal) and TP7-TP8 (temporo-central regions). A vertical and horizontal EOG was recorded from the supra-orbital ridge and the outer canthus of the right eye to monitor blink- and eye-movements for artefact rejection ($>50 \mu V$) in EOG leads. All sites were referenced to the linked ears. The average-reference was recomputed off-line. The EEG signal (impedance < 5kOhm) was amplified (SynAmps: band-pass of 0.1–100 Hz), digitized with 16-bit resolution, sampled at 500 Hz and stored on a hard disk. Epochs (with 100 ms prestimulus baseline) for each tone type were linear detrended, and digitally filtered offline (30 Hz low-pass, 24-dB/octave). ERP data (>50 deviant trials, >600 standard trials) provided a mean of 62-64 MMN sweeps per subject in each group.

Head shapes were digitized by an ultrasound method (Zebris, Munich) to locate skull landmarks (nasion, left/right periauricular points), to determine their positions relative to

the electrodes and improve BESA accuracy. One of the anatomical MR-scans from this study was used for illustrating the results. High resolution T1-weighted 3D anatomical images were recorded with a Siemens 1.5-T sonata with an MP RAGE sequence using a 384 x 384 x 192 matrix giving a 0.5 x 0.5 x 1.0 mm resolution (repetition time/echo time/flip angle = 30 ms/10 ms/30° with a 256 mm field of view).

2.4 Data analysis

Two ERP components were defined by negative-going peaks in the waveform at Fz, FCz and Cz between 90 and 225 ms. The N1 was maximal between 90-120 ms, and the MMN between 100-200 ms in the difference waveform. Separate analyses were run for the N1 and MMN peak amplitudes and latencies. Analyses compared 4 subject groups (EOS, C-EOS, S-15Y, C-15Y) for 29 sites, then for 9 electrodes in the regions of interest at opposite ends of the dipole topography (2 mastoid and 7 fronto-midline recording sites (F3, Fz, F4, FC3, FCz, FC4, Cz: repeat-ed measures) with a repeated measures analysis of variance (ANOVA). The Greenhouse-Geisser epsilon (ϵ) was used as a conservative correction for the degrees of freedom. The locus of effect was determined with one-tailed t-tests for predicted amplitude reductions at frontal and mastoid electrodes.

Detailed analyses with average-referenced data counter the bias towards distant sources of neural activity arising from use of a specific reference: (e.g. use of temporal or mastoid sites in linked-ear referenced data leads to a relative reduction of the potential arising from temporal lobe sites). Mean amplitudes were computed for 5 consecutive time windows (90-105, 105-135, 135-165, 165-195, 105-225 ms). Inspection of the data showed that analyses should concentrate on the 135-165 and 165-195 ms windows covering the MMN peaks. Here, a first set of ANOVAs were run for the 4 subject groups and the 9 electrodes in the regions of interest. A second set of ANOVAs was run using a within-factor of 5 sagittal chains and a within-factor of the 4 electrode sites in these chains (i.e., left; F7, FT7, T3 Ml: left-central; F3, FC3, C3, CP3: central Fz, FCz, Cz, CPz: right-central; F4, FC4, C4, CP4: right; F8, FT8, T4, MR). The aim was to use of a factor with potentially linearly related levels appropriate for the ANOVA and to examine the potential lateral differences for MMN between

groups. Thus post-hoc ANOVAs were restricted to comparisons of the two relevant groups and 3 sagittal chains of electrodes (far left vs. central, far left vs. far right and central vs. far right). A third set of ANOVAs was run on 4 coronal chains and a within factor of 5 electrode sites as in the previous analysis to demonstrate topographic differences between frontal and caudal recordings.

Dipole locations were compared between subjects in any two groups (C-EOS vs. EOS, C-15Y vs. S-15Y, EOS vs. S-15Y, C-EOS vs. C-15Y) with multivariate ANOVAs including the three-level factor location for x, y, and z-coordinates. Similar calculations were made for dipole orientation except that the 2-level factor angle (phi and theta) was included. Group differences of dipole peak strength, and latencies were determined by univariate ANOVAs. Here, a 4-level between-subject factor group (EOS, C-EOS, S-15Y, C-15Y) was included as well as a 4-level within-subject factor dipole (left and right, frontal and temporal lobe locations). Lastly, correlations between medication and ERPs were sought with the Spearman test.

Brain electrical source analysis (BESA, Scherg and Berg 1991) for modelling equivalent dipoles was based on the average-referenced ERP topography where the individual data had been low-pass filtered to avoid aliasing (15 Hz, 24 dB/octave). Dipole computation was based on data from 20 ms before to 40 ms after the peak. BESA allows iterative spatio-temporal fitting of the dipole location and orientation in a spherical head-model with an inner radius of 85 mm corrected for brain, skull, and scalp conductivity (Scherg and Picton 1991), until the difference between the recorded surface data and the calculated surface data of the dipole-model is minimised (least square fit). The goodness of fit is expressed by the mean residual variance (RV) and the best fit is the lowest RV not explained by the model, as a percentage of the signal variance within the given time window.

The dipole model, were led by the hypothesis of Jemel et al. (2002), required bilateral frontal and temporal lobe dipoles to explain >95% of the topographic variance. This 4-dipole solution has proved robust and replicable (Oknina et al., 2004). The model was first fitted to the ERP grand mean, that describes the spatio-temporal data

matrix, and then applied to the individual subjects' data. During the iterative best-fitting estimations the temporal followed by the frontal lobe dipoles were initially constrained to the hypothesis, and then allowed to move without constraint. The final unconstrained group solution had an RV < 2%, and matched that of Jemel et al (2002).

Stereotaxic Talairach coordinates for these sources were calculated (Garnero et al. in Crottaz-Herbette and Ragot, 2000), and 2 methods used to represent the locations on brain images. First, the group solution was mapped on to the axial, coronal and sagittal brain MR-images of a healthy subject from the C-15Y group. Then, individual source solutions were calculated. To account for the decreased signal-to-noise ratio of a single subject's waveform, individual solutions were re-calculated to a criterion of RV < 4%. The distances of the individual best solutions from the mean of the individual solutions was mapped on to the brain image of an adult control normalised with reference to the SPM99 brain template with the radius of the circle representing the standard deviation from the mean locations of the individual solutions.

3. Results

3.1 ERP Analysis

In Fig. 1 the deviant and standard N1 components and the MMN at FCz (average-reference) are compared for each group. There were group differences for the N1 peak amplitude for both types of stimulus [$F(24, 536) = 3.3$ - 5.7 , $p < 0.001$, $\epsilon = 0.21$ - 0.18]. In EOS patients, N1 amplitudes after both tones [$F(8, 296) = 7.2$ - 13.1 , $p < 0.003$ - 0.001 , $\epsilon = 0.23$ - 0.20] were smaller than in controls at all fronto-central electrodes ($t_{37} > 2.3$, $p < 0.025$: standards -3.7 vs. -6.2 μV). N1 attenuation in the S-15Y vs. the C-15Y group was restricted to the standard tones [$F(8, 240) = 4.67$, $p < 0.03$, $\epsilon = 0.16$] at Fc3 and Cz [$t(30) > 2.58$, $p < 0.05$: -4.0 vs. -5.6 μV].

ANOVA for the mean MMN amplitudes of the average-referenced data (135-165 ms) for the 9 electrode sites of interest revealed significant group by electrode interactions [$F(24, 536) = 2.2$, $p = 0.048$, $\epsilon = 0.25$]. In the following window group differences were significant after covarying for IQ [$F(24, 536) = 1.5$, $p = 0.19$: $F(24, 424) = 2.1$,

$p = 0.04$, $\epsilon = 0.33$]. Each patient group was then compared with their age-matched controls. In the younger age-class an indication for a group by electrode interaction [$F(8, 296) = 2.55$, $p = 0.08$, $\epsilon = 0.27$: 135-165 ms] was explained by a small frontal MMN amplitude decrease [$t(37) = 2.0$, $p < 0.027$: FC4] and a marked 50% reduction in the mastoid recording from EOS patients [$t(37) = -2.7$, $p < 0.006$] that continued over the next 30 ms time window [$t(37) = -2.2$, $p < 0.018$: M2]. A modest interaction for the 2 older groups, especially in the 165-195 ms window [$F(8, 240) = 3.0$, $p = 0.06$, $\epsilon = 0.23$] was more marked after covarying for IQ [$F(8, 208) = 5.2$, $p = 0.01$, $\epsilon = 0.23$]. Unlike the younger patients, the interaction for S-15Y patients was accounted for by an amplitude reduction at the left mastoid [135-165 ms, $t(30) = -1.9$, $p = 0.031$] not at frontal sites. This effect was marked at both mastoids in the 165-195 ms window [$t(30) = -2.5$, $p = 0.01$: M1]. A comparison of the younger with the older patients showed no improvement or progression of the impairment (135-195 ms, $F = 1.5$ - 0.9). But, a trend for a further decrease of amplitude at the left mastoid by 1 μV with age was observed [135-165 ms, unprotected $t(40) = 2.1$, $p = 0.022$: Table 1].

A topographic change was supported by the ANOVA comparing coronal lines of electrodes in both time windows ($F(36, 804) = 1.78$ - 1.94 , $p = 0.037$ - 0.06 , $\epsilon = 0.28$ - 0.30). Significant change was restricted to the younger subjects, and showed that the normal fronto-caudal difference was reduced in the patients (e.g. electrode line 2 vs. 4, $F(4, 148) = 3.8$, $p = 0.04$, $\epsilon = 0.58$). Analysis of the sagittal lines of electrodes for laterality effects showed an overall trend [$F(36, 804) = 1.8$, $p = 0.065$, $\epsilon = 0.28$: 135-165 ms]. Modest changes in the EOS vs. C-EOS group [$F(12, 444) = 2.3$, $p = 0.066$, $\epsilon = 0.30$] were explained by the central vs. right [$F(3, 111) = 4.8$, $p = 0.015$, $\epsilon = 0.56$] and left vs. right comparisons [$F(3, 111) = 2.7$, $p = 0.08$, $\epsilon = 0.56$]. The reduction was pronounced at the right mastoid ($t(37) = -2.68$, $p = 0.005$). Comparisons over 165-195 ms continued to indicate a difference [$F(12, 444) = 2.6$, $p = 0.05$, $\epsilon = 0.27$] with a right lateralized MMN reduction [central vs. right: $F(3, 111) = 6.4$, $p = 0.005$, $\epsilon = 0.54$] being marked at the mastoid [$t(37) = -2.19$, $p = 0.018$].

Figure 1

The MMN (below) and the deviant/standard N1 waveforms (above) are shown (average-reference) for 4 subject groups. Mean peak MMN latencies at FCz did not differ between patients and their respective controls. MMN amplitudes (see table 1) were significantly smaller in the patients than the controls, especially at the mastoids at and after attaining peak amplitude (in the windows covering 135-195 ms). See text and table 1 for details about the N1.

In summary, these analyses lead to two predictions on source activity. First patients may show but modest differences at frontal sites (temporal lobe generators?). Second differences appear more marked at mastoid sites, on the right in the younger and less clearly on the left in the older patients. The differences could reflect the activity of frontal generators.

3.2 Dipole Source Analysis

Dipole source locations for 60 ms of activity around the MMN peak were determined for each group (Fig. 2). The waveforms of all groups were well explained by the 4-dipole model (Jemel et al 2002). Minor changes to the dipoles' location and orientation led to residual variances (RV) below 1.5 %, and best fits

Table 2

Localisation, Strength and Latency of Modelled Dipole Sources in Younger and Older Patients with Schizophrenia and their Age-matched Healthy Controls: On the Left for the Group Solution, on the Right for the Mean of the Individual Solutions

Subject Group (N)	Time Window (ms)	Residual Variance (best fit %)	x	y	z	Hemi-sphere	BA	Brain Region	Strength nA	Latency ms	N	x (SD)	y	z	Strength nA (SD)	Latency ms (SD)
C-15Y (15)	90-150	0.90 (0.43)	-56 ²	-23	+10	Left	41	STG	18.2	120	12	-53 (5)	-29 (5)	11 (5)	25.8 (18.6)	126 (49)
			+59	-15	+1	Right	22	STG	15.6	128		56 (6)	-17 (4)	5 (4)	25.1 (24.4)	124 (50)
			-8	-14 ³	+42	Left	24	ACG	10.5	112		-8 (4)	-18 (3)	37 (4)	20.4 (16.7)	120 (50)
			+36	+45	+19 ⁴	Right	10	MFG	9.8	134		34 (5)	40 (3)	17 (4)	18.3 (21.4)	134 (52)
S-15Y (17)	96-156	1.05 (0.75)	-46 ²	-28	+9	Left	41	STG	16.6	124	14	-45 (5)	-28 (8)	8 (6)	20.3 (7.2)	123 (16)
			+52	-15	-2 ⁷	Right	22	STG	19.3	116		53 (6)	-19 (4)	1 (7)	17.2 (6.2)	133 (23)
			-7	+2 ^{3,6}	+39	Left	24	ACG	10.0	124		-8 (5)	-4 (5)	36 (5)	16.0 (8.3)	125 (23)
			+38	+43 ⁵	+8 ⁴	Right	10	MFG	12.0	154		39 (5)	38 (4)	8 (6)	12.7 (6.3)	139 (27)
C-EOS (14)	110-170	1.24 (0.62)	-44	-36	+9	Left	41	STG	28.9	126	11	-47 (5)	-32 (6)	9 (6)	32.1 (10.2)	136 (26)
			+55	-11	-0 ¹	Right	22	STG	23.0	116		58 (4)	-14 (6)	5 (5)	19.5 (8.0)	141 (18)
			-3	-10	+31	Left	24	ACG	14.6	128		-7 (5)	-13 (6)	33 (7)	31.7 (15.9)	137 (31)
			+42	+46	+11	Right	10	MFG	8.1	158		42 (6)	40 (4)	9 (5)	18.8 (7.5)	144 (30)
EOS (25)	96-156	1.45 (0.66)	-44	-29	+6	Left	41	STG	20.9	122	18	-45 (7)	-28 (8)	6 (6)	27.0 (9.7)	139 (32)
			+54	-14	+12 ^{1,7}	Right	41	S/TTG	12.4	144		53 (7)	-17 (6)	10 (6)	22.9 (11.1)	142 (36)
			-10	-6 ⁶	+42	Left	24	ACG	8.8	140		-9 (5)	-8 (7)	41 (7)	23.1 (12.1)	135 (37)
			+41	+46 ⁵	+10	Right	10	MFG	8.4	130		39 (5)	43 (5)	8 (6)	14.3 (7.3)	151 (33)

ACG, anterior cingulate gyrus; BA, Brodmann area; MFG, Mid-frontal gyrus; STG, superior temporal gyrus; TTG, transverse temporal gyrus

In the young patients (vs. young controls) the right temporal source tended to be more dorsal¹: In the older patients (vs. older controls) the left temporal source was more medial², the left cingulate source was rostral³, and the right frontal source more ventral⁴: Compared to young patients, dipole sources in the older patients tended to be more caudal in the right frontal lobe⁵, but continue to be more rostral in the left cingulate⁶, while in the right temporal lobe the source tended to return to the more ventral locus⁷ seen in controls. For further details see the text.

Figure 2

Spatio-temporal multiple dipole analysis of the MMN component for the grand-mean ERP of each of the S-15Y and C-15Y groups on the left, and the EOS and C-EOS group on the right. For each group there is the temporal development of each of the 4 dipole moments and the residual variance (ca. 1%) for the model on the left: vertical bars show the 60 ms analysis period. The stereotaxic coordinates for their locations are shown in sections from the brain atlas in the centre and right (Talairach and Tournoux, 1988). They show two mildly asymmetrical bilateral temporal lobe dipoles (BA 41, 42) and two strongly asymmetrical bilateral frontal lobe dipoles (BA 24, 10)

below 0.8 % for all groups. Table 2 (left) summarises the coordinates for all 4 group solutions, the RV of activity not explained by the model, the best fits in each group, the dipole moments, and their latencies. The two posterior dipoles lay bilaterally in the superior temporal gyri (BA 41/BA 22) with that in the left hemisphere more dorsal and posterior than the one in the right hemisphere. The frontal dipoles were located posterior in the left anterior cingulate gyrus (BA 24) and in the right inferior/mid-frontal gyrus (BA 10). Their locations and orientations are illustrated in Figure 3. The group solutions can be seen to differ slightly from each other.

3.3 Group Comparisons of Dipole Location

To calculate statistical differences, the respective group solution was fitted to the individual's MMN. For most subjects (Table 2) only minor changes in orientation and location were needed to reach $RV < 4\%$. The results (the locus diameter represents 2 SD) are mapped on to a control brain (Fig. 4). The changes along a given dimension in millimetres reflect the range of values from

the BESA group mean [$RV < 2\%$] and the mean of the individual solutions [$RV < 4\%$: table 2].

Compared to the young controls, the EOS patients' right temporal lobe source [Wilks $\Lambda = 0.55$, $F(3, 25) = 6.7$, $p = 0.032$] was about 6-12 mm more dorsal [$F(1, 27) = 6.1$, $p < 0.02$]. However, between adult patients and their controls differences were marked for both frontal dipoles [$\Lambda_s = 0.22-0.39$, $F_s(3, 22) = 11.5-26.6$, $p < 0.002$]. For adult patients the left cingulate source was more anterior [16 mm; $F(1, 24) = 68.5$, $p < 0.002$] and the right frontal source was more ventral [c. 8-11 mm; $F(1, 24) = 16.9$, $p < 0.002$]. In addition left temporal lobe sources tended to be 8-10 mm more medial nearer to the locus noted in young subjects, above [$F(1, 24) = 14.9$, $p < 0.02$].

EOS and S-15Y patients were compared for potential signs of illness progression with duration. In S-15Y patients the right frontal locus was ca. 5 mm more caudal [$\Lambda = 0.72$, $F(3, 28) = 3.7$, $p < 0.05$; $F(1, 30) = 11.0$, $p = 0.032$]. The left cingulate source, already more rostral

Figure 3

Source locations for the two frontal dipoles based on BESA group mean solutions for healthy adult controls (C-15Y) and the older patient group (S-15Y) have been mapped on to an image of the brain of a healthy subject in the C-15Y group. The more anterior location of the left cingulate and the less dorsal location of the right inferior/mid-frontal sources in the patient group are clearly illustrated.

in the EOS than in the controls, was yet a further 3-7 mm more rostral [$\lambda = 0.62$, $F(3, 28) = 5.7$, $p < 0.05$]. In the right temporal lobe the source tended to return 9-14 mm from a more dorsal (in the EOS) to a more usual and more ventral locus seen in other groups [$\lambda = 0.63$, $F(3, 28) = 5.5$, $p = 0.064$; $F(1, 30) = 14.4$, $p < 0.02$].

There were no differences between groups for the peak latencies of the dipole moments ($F < 1$). This lack may be attributed to the variability of the individual differences (Table 2). However, it is important to note that independent of subject group the right mid-frontal source peaked significantly later (143 ms) than that in the left cingulate (130 ms) or in the left temp-oral lobe (131 ms) illustrating a bottom-up sequence of activation [$F(3, 153) = 5.7$, $p < 0.01$; $t(54) > -3.6$, $p < 0.05$].

3.4 Dipole Strength and Orientation

Across groups the peak dipole strengths differed significantly from each other [$F(3, 153) = 16.9$, $p < 0.01$]. The lowest activity in the right frontal source [15.6 nA; $t(54) > 4.7$, $p < 0.05$] contrasted with sources in the temporal lobes (left: 26.1 nA, right: 21.2 nA) and the left cingulate gyrus (22.4 nA). Further, the peak dipole strength was higher in the left than the right temporal lobe [$t(54) = 3.1$, $p = 0.05$]. Differences between groups were small [$F(9, 153) = 2.3$, $p = 0.028$], but dipole strength in the left temporal lobe of the S-15Y group (20.3 nA) was weaker than in the young first-episode patients (27.0 nA; $t(30) = 2.2$, $p = 0.039$], consistent with some progression of the disorder. Lateralization ratios (left-minus-right/left-plus-right) confirmed that while there was a leftward asymmetry for temporal lobe dipole moment in healthy subjects, this asymmetry was absent in the young patients [EOS: 0.096 SD 0.25 vs. C-EOS: 0.248 SD 0.19; $t(27) = 1.75$, $p < 0.046$, 1-tail]. Asymmetries and differences were not significant for the older subjects or for the frontal dipole moments.

Analyses of the dipoles' orientations yielded no significant patient-control differences. This likely reflected the large variation between individual values that themselves were calculated on the basis of poor signal-to-noise ratio ratios in the few trials available (Table 2). But a trend was

noted for the cingulate dipole [$\lambda = 0.67$, $F(2, 23) = 5.7$, $p = 0.01$] to be left-ventrally oriented in the EOS group and right-ventrally oriented in the S-15Y patients [$F(1, 24) = 1.0$, $p = 0.003$].

3.5 Medication:

N1 amplitudes after the standard and deviant tones did not differ between the 13 EOS patients on medication and the 11 tested without ($F < 1.5$). MMN amplitudes were similar for those on (Fz: $-4.8 \mu V$ SD 1.7) and those off medication (Fz: $-4.2 \mu V$ SD 2.9; $F < 1.0$). Shorter latencies for medicated vs. unmedicated patients did not differ significantly (124 ms SD 20 vs. 156 ms SD 37). There were no amplitude differences in the window around the MMN peak (135-165 ms; $F < 1.0$). But in the following window (165-195 ms) the morphology of the waveform showed a slower fall-off from the negative peak toward baseline in the non-medicated patients. This resulted in their MMN being larger over this later period (-0.9 vs. $0.0 \mu V$: $F(8, 176) = 4.8$, $p = 0.016$, $\epsilon = 0.23$).

4. Discussion

The main purpose of this study was to explore ways to measure, calculate and present four features of the frequency-deviant MMN dipole sources (dipole moment, peak latency, brain location and orientation) active in auditory change detection and to relate these to the processes of psychopathology and illness progression.

There was a modest reduction of MMN amplitude in adolescent patients with early onset schizophrenia. This was qualitatively like previous reports (e.g. Catts et al., 1995; Javitt et al., 1995). The reduction was less than we reported previously (Oades et al., 1995; 1996), but similar to or more than in other recent reports (Bramon et al., 2004; Michie et al., 2000). The decrease was slightly more marked in the acutely ill EOS than the S-15Y patients first diagnosed 15 years before. It was most evident at the mastoid sites where a two thirds reduction from the peak latency for the following 60 ms was recorded. Frontal N1 amplitudes elicited by the tones were also reduced in both patient groups.

Figure 4

Estimates of the group centre of activation and the spread of the individual's best fit are shown for all 4 groups. Dipole-sources are represented by the means of the solutions in each of the 3 dimensions for those subjects for whom the mean group solution was adapted to account for at least 96% of the variance in the individual's data. The radius of each location in any dimension represents the SD of the mean of the individual solutions: the data are shown in table 2 on the right.

This difference occurred 60+ ms before the reduction of MMN in patients. Both features militate against a major role for N1 alterations in the reduction of MMN. A decreased N1 amplitude is consistent with some patient samples (Alain et al., 2002; Ford et al., 1994), but not others (Shinozaki et al., 2002). Medication in 13/24 EOS patients produced a sharper MMN peak that contrasted with a longer lasting MMN morphology in the non-medicated patients.

The dipole source locations broadly replicated our previous report (Jemel et al., 2002) and are generally consistent with the bilateral temporal and frontal sources recently described using the minimum norm method (Valkonen-Korhonen et al., 2003). But here the sources already tended to be more dorsal

in the right temporal lobe of adolescent EOS-patients compared to age-matched controls. The left frontal source, posterior in the anterior cingulate, tended to be more rostral in the young patients and significantly so in the adult S-15Y group. Further signs of structural change, perhaps reflecting illness progression, were also evident in the right mid-frontal source location that lay more caudal in the adult patients compared to the younger patients and controls.

4.1 Schizophrenia:

Impairments of MMN have been reported in adolescent (Oades et al., 1997) and adult patients with schizophrenia of recent onset (Shinozaki et al., 2002). But others report that the impairment may take some time to develop (Grzella et al., 2001; Umbricht et al.,

2002; Salisbury et al., 2002). For the first time we report here putative changes in the bases generating MMN that may be responsible for the widespread reports of reduced MMN in schizophrenia (see introduction).

In all cases 2 sources lay in the superior temporal gyrus, probably summing the activity in primary and secondary auditory cortices (Kircher et al., 2004). The locus on the right was antero-lateral to that on the left, as reported for the MEG equivalent of MMN (Kasai et al., 2003). [But note that relative to the primary auditory cortex the source lay more posterior in secondary association areas.] In general sources were stronger on the left, but this asymmetry was not found in the patients. For comparison, Kircher et al. (2004) reported a rightward asymmetry for duration-deviant MMN in controls that was also absent in their chronically ill patients. Here, in younger patients the right temporal source was more dorsal than in controls. In contrast, in older patients the anomaly lay on the left, where the temporal lobe source was located more medially. There is good evidence that the function of temporal lobe generators relates to the maintenance of a working memory-like trace of the usual auditory stimulation (Berti and Schröger, 2003): event-related imaging has shown that such information can access directly the inferior frontal cortex (D'Esposito et al., 2000). Indeed, a cingulate contribution to the maintenance of auditory memory has also been described (Shibata and Yukie, 2003). Further, anatomical tracing studies have shown that both auditory association areas and the cingulate region send input to the inferior frontal gyrus (Petrides and Pandya, 2001).

The left anterior cingulate source in both patient groups lay anterior to their respective controls. The right mid-frontal source was more ventral in the older patients than in their respective controls. It is interesting that an fMRI investigation of musicians' response to inconsonant chords showed activity specific to similar regions (Tillman et al., 2003), including the inferior/ mid-frontal gyri, the insula, and the bilateral cingulate cortex. Patients with schizophrenia have been reported to have smaller inferior frontal volumes, which correlated with mid-frontal volumes

(Buchanan et al., 2004). Activity in this region has been noted in other paradigms that, like the MMN condition, require pre-conscious attention-related processing (e.g. back-ward masking, sensory gating), and to be reduced in patients with schizophrenia (Dehaene et al., 2004; Kumari et al., 2003).

Generic detection of stimulus change has been attributed to right inferior mid-frontal function (Beck et al., 2001). This conclusion was based on the activity elicited in an fMRI investigation of change-blindness during the condition of conscious non-detection of actual stimulus change. Such frontal pre-conscious activity is proposed to represent the switch to further controlled processing of a potentially significant stimulus (Escera et al., 1998). Numerous paradigms have postulated that parts of the cingulate cortex are involved in monitoring conflicts of information processing (e.g. error commission) precisely as would occur in the current mismatch conditions. This is illustrated by Carter et al. (2001) who used a form of Go/no-go task. They also reported activity in the right frontal BA 10 region. Our finding that dipole activity in the right frontal region, occurred later than the near-simultaneous activation of the auditory association cortices and cingulate, provides remarkable support for a bottom-up sequence of processing and monitoring that culminates in a region exerting a "pre-executive" role necessary for subsequent controlled processing.

4.2 Illness Progression:

Controversy surrounds whether there is a 'progression of illness' in patients with schizophrenia over time in terms of function or structure. Potential signs of structural change were the more rostral position of the left cingulate and the more ventro-caudal location of the right frontal MMN sources in older vs. younger patients. The relevance is underlined by the activity of the right frontal dipole peaking precisely in the time window for which a significant difference in the ERP was recorded. The change in the cingulate locus was also accompanied by an altered orientation of the dipole between younger and older patients.

Modest structural and functional

progression was also observed for the left temporal sources. The locus remained more medial rather than developing a more lateral position seen in age-matched controls. Further, the dipole moment was significantly weaker in the older vs. the younger patients. A similar emphasis on left sided impairments was reported for MMN in adult patients (Catts et al 1995; Hirayasu et al 1998; Kasai et al 2003; Sauer et al 1997). Our results are consistent with a fairly marked change of structure for patients in adolescence (Gogtay et al., 2004) that asymptotes over the following decade (DeLisi et al., 2004). The later changes in the superior temporal gyrus and frontal regions were reported to become more pervasive with age (Thompson et al., 2001), and could be indirectly responsible for the changes of location for the generators described here. Such changes with increasing age reflect differences in grey to white matter ratios driven by the absence of white matter expansion in patients (Bartzikis et al., 2003). Changes of function with the course of illness may incur executive improvements, yet memorial impairments (Hoff et al., 1999). Theoretically both domains of function could relate to appropriate MMN generation.

4.3 Limits to the Study:

A number of features qualify the accuracy of dipole localization. The first concerns the variance in the data. The two patient groups do not represent subjects with a homogeneous set of states or traits that are likely to provide a uniform set of MMN measures or dipole locations. Thus, the EOS group contains subjects with a paranoid, disorganized or undifferentiated form of illness. Clearly the presence or absence of paranoia or thought disorder will influence, at least quantitatively, the potentials recorded (Oades et al., 1996; Krljes et al., 2004). Similarly S-15Y patients manifested varying degrees of illness-severity. Further, some of the patients were not medicated at the time of the recording. These provide further sources of variance. While it is reported that the dopaminergic aspect of anti-psychotic medication does not have a major influence on MMN expression (Käkonen et al., 2000; Hansenne et al., 2003), other aspects of the drug profile have not been studied. While

medication did not significantly affect peak amplitude or latency here, it did prolong the period of MMN activation. The group participants were not precisely matched for gender, but studies of the potential influence of gender on MMN have been consistently negative (Koelsch et al 2003; Nagy et al 2003). These aspects will be taken into account in the ongoing longitudinal design of the study.

A second set of considerations concerns the signal to noise ratio in the source calculations. A reasonable effect size requires about 1000 trials of artefact free data derived from 6 electrodes for each assumed dipole source (Kamijo et al., 2002; Phillips et al., 2002; Picton et al., 1995). Each group solution satisfied the data requirement in terms of the number of sweeps and channels with a small excess of degrees of freedom. But this would not allow the additional resolution of, say, 2 dipoles in each temporal lobe. In fact, our coordinates overlap with those reported for to N1/P2 dipoles attributed to the primary auditory cortex (Mulert et al., 2002). The stability of the group solution was aided by the hypothesis-led model (Jemel et al., 2002). But the individual solutions, calculated to provide a measure of variance, were less reliable due to a much poorer signal-to-noise ratio. We had to relax the criterion for the model's fit and stability. Nonetheless a satisfactory 70-80% of the subjects achieved the criterion of explaining 96% of the variance in the MMN recorded.

A final set of considerations lies with the spatial accuracy of the solutions. While topographic precision is improved by digitizing the recording sites with respect to individual fiducial skull markers, we must emphasize that the model to explain these data provides a mathematical point solution. This does not plausibly reflect the likely extent of physiological activation. We did not check the stability of solutions millimetre by millimetre along the 3 dimensions, as before (Jemel et al., 2003). But the stability we reported (2-6 mm along each dimension) is likely to pertain here where all other aspects of our methods were the same. Further, the standard deviations (table 2) show that this measure of variation was usually <8 mm. However, these difficulties are compounded by the

inexactitude of placing sources on the brain atlas or SPM maps based on the anatomy derived from averages of other subjects. The problem is partly counteracted by our placing sources on the anatomy of one of our subjects (fig. 3), but this does not account for anatomical variability within our subjects. In the future, the data should be placed on anatomic averages of the subjects studied. Currently, in conjunction with studies of the accuracy obtained with known modelled sources (e.g. Yvert et al 1997; Laarne et al 2000), we postulate a potential error of the order of 1 cm.

In conclusion we have shown that our model describing 4 dipole sources for the frequency-deviant MMN (Jemel et al., 2002) broadly fits groups of healthy adolescent and adult subjects and age-matched patients with schizophrenia. These sources lie bilaterally in the auditory association cortices, in the posterior part of the left anterior cingulate and the right inferior or mid-frontal gyrus. Some changes in these source locations are related to pathophysiology and illness progression. The reduction of MMN to be found in many patients with schizophrenia is reflected in small changes to each source, differentially reflecting the size of the dipole moment, its orientation and its location.

References

- Alain, C., Cortese, F., Bernstein, L.J., He, Y., Zipursky, R.B., 2002. Auditory feature conjunction in patients with schizophrenia. *Schizophr Res.* 49, 179-191.
- Arrington, C. M., Carr, T. H., Mayer, A. R., Rao, S. M., 2000. Neural mechanisms of visual attention: object-based selection of a region in space. *J. Cogn Neurosci.* 12, S106-117.
- Bartzokis, G., Nuechterlein, K.H., Gitlin, M., Rogers, S., Mintz, J., 2003. Dysregulated brain development in adult men with schizophrenia: a magnetic resonance imaging study. *Biol. Psychiatry* 53, 412-421.
- Beck, D.M., Rees, G., Frith, C.D., Lavie, N., 2001. Neural correlates of change detection and change blindness. *Nature Neurosci.* 4, 645-650.
- Benkert, O., Hippus, H., 1986. *Psychiatrische Pharmakotherapie* (4th ed.). Heidelberg: Springer-Verlag
- Berti, S., Schröger, E., 2003. Working memory controls involuntary attention switching: evidence from an auditory distraction paradigm. *Eur. J. Neurosci.* 15, 1119-1122.
- Bramon, E., Croft, R.J., McDonald, C., Virdi, G.K., Gruzelier, J.G., Baldeweg, T. et al., 2004. Mismatch negativity in schizophrenia: a family study. *Schizophr. Res.* 67, 1-10.
- Brazdil, M., Roman, R., Falkenstein, M., Daniel, P., Jurak, P., Rektor, I., 2002. Error processing-evidence from intra-cerebral ERP recordings. *Exp Brain Res* 146:460-466.
- Buchanan, R.W., Francis, A., Arango, C., Miller, K., Lefkowitz, D.M., McMahon, R.P. et al., 2004. Morphometric assessment of the heteromodal association cortex in schizophrenia. *Am. J. Psychiatry* 161, 322-331.
- Catts, S.V., Shelley, A-M., Ward, P.B., Liebert, B., McConaghy, N., Andrews, S., Michie, P.T. 1995. Brain potential evidence for an auditory sensory memory deficit in schizophrenia. *Am J Psychiatry*, 152, 213-219.
- Carter, C.S., MacDonald, A.W., Ross, L.L., Stenger, V.A., 2001. Anterior cingulate cortex activity and impaired self-monitoring of performance in patients with schizophrenia: an event-related fMRI study. *Am. J. Psychiatry* 158, 1423-1428.
- Carter, C.S., Braver, T.S., Barch, D.M., Botvinick, M.W., Noll, D., Cohen, J.D., 1998. Anterior cingulate cortex, error detection and the online monitoring of performance. *Science* 280, 274-279.
- Crottaz-Herbette, S., Ragot, R., 2000. Perception of complex sounds: N1 latency codes pitch and topography codes spectra. *Electroencephalogr. Clin. Neurophysiol.* 111, 1759-1766.
- Dehaene, S., Artiges, E., Naccache, C., Martelli, C., Viard, A., Schürhoff, F. et al., 2004. Conscious and subliminal conflicts in normal subjects and patients with schizophrenia: the role of the anterior cingulate. *Proc. Natl. Acad. Sci. USA* 100, 13722-13727.
- Delisi, L.E., Maurizio, A., Sakuma, M., Hoff, A.L., 2004. Progressive ventricular enlargement: a 10-year follow-up study. *Schizophr. Res.* 67, 27.
- D'Esposito, M., Postle, B.R., Rypma, B., 2000.

- Prefrontal cortical contributions of working memory: evidence from event-related fMRI studies. *Exp. Brain Res.* 133, 3-11.
- Eggers, C., Bunk, D., Volberg, G., Röpcke, B., (1999). The ESSEN study of childhood onset schizophrenia: Selected results. *Eur. Child Adolesc. Psychiatry* 8: suppl.1, 21-28.
- Escera, C., Alho, K., Winkler, I., Näätänen, R., 1998. Neural mechanisms of involuntary attention to acoustic novelty and change. *J. Cogn. Neurosci.* 10, 590-604.
- Erikson, R., Goldthorpe, J.H., 1992. Individual or family? Results from two approaches to class assignment. *Acta Sociol.* 35, 95-106.
- Ford, J.M., White, P.M., Csernansky, J.G., Faustman, W.O., Roth, W.T. et al., 1994. ERPs in schizophrenia: effects of antipsychotic medication. *Biol. Psychiatry* 36, 153-170.
- Gogtay, N., Sporn, A., Clasen, L.S., Nugent, T.F., Greenstein, D., Nicolson, R. et al., 2004. Comparison of progressive cortical gray matter loss in childhood-onset schizophrenia with that in childhood-onset atypical psychoses. *Arch. Gen. Psychiatry* 61, 17-22.
- Grzella, I., Müller, B.W., Oades, R.D., Bender, S., Schall, U. et al., 2001. Novelty-elicited mismatch negativity (MMN) in patients with schizophrenia on admission and discharge. *J. Psychiatry. Neurosci.* 26, 235-246.
- Hansenne, M., Pinto, E., Scantamburlo, G., Couvreur, A., Reggers, J., Fuchs, S. et al., 2003. Mismatch negativity is not correlated with neuroendocrine indicators of catecholaminergic activity in healthy subjects. *Hum. Psychopharmacol. Exp. Clin.* 18, 201-205.
- Harris, L.R., Lieberman, L. 1996. Auditory stimulus detection is not suppressed during saccadic eye movements. *Perception* 25, 999-1004.
- Hirayasu, Y., Potts, G.F., O'Donnell, B.F., Kwon, J.S., Arakaki, H., Akdag, S.J. et al., 1998. Auditory mismatch negativity in schizophrenia: topographic evaluation with a high-density recording montage. *Am. J. Psychiatry* 155, 1281-1284.
- Hoff, A.L., Sakuma, M., Wieneke, M., Horon, R., Kushner, M., Delisi, L.E., 1999. Longitudinal neuropsychological follow-up study of patients with first-episode schizophrenia. *Am. J. Psychiatry* 156, 1336-1341.
- Jacobsen, T., Schröger, E., 2001. Is there pre-attentive memory-based comparison of pitch? *Psychophysiology* 38, 723-727.
- Javitt, D.C., Doneshka, P., Grochowski, S., Ritter, W., 1995. Impaired mismatch negativity generation reflects widespread dysfunction of working memory in schizophrenia. *Arch. Gen. Psychiatry* 52, 550-558.
- Jemel, B., Achenbach, C., Müller, B., Röpcke, B., Oades, R.D., 2002. Mismatch negativity results from bilateral asymmetric dipole sources in the frontal and temporal lobes. *Brain Topogr.* 15, 13-27.
- Jemel, B., Oades, R.D., Oknina, L., Achenbach, C., Röpcke, B., 2003. Frontal and temporal lobe sources for a marker of controlled auditory attention: the negative difference (Nd) event-related potential. *Brain Topogr.* 15, 249-262.
- Kähkönen, S., Ahveninen, J., Pekkonen, E., Kaakkola, S., Huttunen, J., Ilmoniemi, R.J., Jääskeläinen, I.P., 2002. Dopamine modulates involuntary attention shifting and reorienting: an electromagnetic study. *Clin. Neurophysiol.* 113, 1894-1902.
- Kamijo, K-I., Yamazaki, T., Kiyuna, T., Takaki, Y., Kuroiwa, Y., 2002. Visual event-related potentials during movement imagery and the dipole analysis. *Brain Topogr.* 14, 279-292.
- Kane, J.M., 1996. Drug therapy: Schizophrenia. *New England J. Med.* 334, 34-41.
- Kasai, K., Yamada, H., Kamio, S., Nakagome, K., Iwanami, A., Fukuda, M. et al., 2003. Neuromagnetic correlates of impaired automatic categorical perception of speech sounds in schizophrenia. *Schizophr. Res.* 59, 159-172.
- Kircher, T.T.J., Rapp, A., Grodd, W., Buchkremer, G., Weiskopf, N., Lutzenberger, W. et al., 2004. Mismatch negativity responses in schizophrenia: a combined fMRI and whole-head MEG study. *Am. J. Psychiatry* 161, 294-304.
- Krljes, S., Hirsch, S.R., Baldeweg, T., 2004. Event-related potential mismatch negativity is related to thought disorder in schizophrenic patients. *Schizophr. Res.* 67, 22.

- Koelsch, S., Maess, B., Grossmann, T., Firederici, A.D., 2003. Electric brain responses reveal ender differences in music processing. *NeuroReport* 14, 709-713.
- Kumari, V., Gray, J.A., Geyer, M.A., ffytche, D., Soni, W., Mitterschiffthaler, M.T. et al., 2003: Neural correlates of tactile prepulse inhibition: a functional MRI study in normal and schizophrenic subjects. *Psychiatry Res. Neuroimaging* 122, 99-113.
- Laarne, P.H., Tenhunen-Eskelinen, M.L., Hyttinen, J.K., Eskola, H.J., 2000. Effect of EEG electrode density on dipole localization accuracy using two realistically shaped skull resistivity models. *Brain Topogr.* 12, 249-254.
- Michie, P.T., Budd, T.W., Todd, J., Rock, D., Wichmann, H., Box, J. et al., 2000. Duration and frequency mismatch negativity in schizophrenia. *Clin. Neurophysiol.* 111, 1054-1065.
- Mulert, C., Juckel, G., Augustin, H., Hegerl, U., 2002. Comparison between the analysis of the loudness dependency of the auditory N1/P2 component with LORETA and dipole source analysis in the prediction of treatment response to the selective serotonin reuptake inhibitor citalopram in major depression. *Clin. Neurophysiol.* 113, 1566-1572.
- Näätänen, R., Gaillard, A.W.K., Mäntysalo, S., 1978. Early selective attention effect on evoked potential reinterpreted. *Acta Psychol.* 42, 313-329.
- Nagy, E., Potts, G.F., Loveland, K.A., 2003. Sex-related differences in deviance detection. *Int. J. Psychophysiol.* 48, 285-292.
- Oades, R.D., 1995. Connections between studies of the neurobiology of attention, psychotic processes and event-related potentials. *Electroencephalogr. Clin. Neurophysiol. Suppl.* 44, 428-438.
- Oades, R.D., Zerbin, D., Dittmann-Balcar, A., Eggers, C., 1996. Auditory event-related potential (ERP) and difference-wave topography in schizophrenic patients with/without active hallucinations and delusions: a comparison with young OCD and healthy subjects. *Int. J. Psychophysiol.* 22, 185-214.
- Oades, R.D., Dittmann-Balcar, A., Zerbin, D., Grzella, I. 1997. Impaired attention-dependent augmentation of MMN in nonparanoid vs. paranoid schizophrenic patients: a comparison with obsessive-compulsive disorder and healthy subjects. *Biol. Psychiatry* 41, 1196-1210.
- Ofek, E., Pratt, H., 2004. Ear advantage and attention: an ERP study of auditory cued attention. *Hearing Res.* 189, 107-118.
- Oknina, L.B., Juran, S.A., Oades, R.D., Weisbrod, M., Chan, E., Chen, E.Y.H. et al., 2004. Frontal and temporal lobe sources for mismatch negativity (MMN) in schizophrenia: an ERP and MR-anatomical imaging study. *Schizophr. Res.* 67, 22.
- Oldfield, R.C., 1971. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9, 97-113.
- Paavilainen, P., Alho, K., Reinikainen, K., Sams, M., Näätänen, R., 1991. Right hemisphere dominance of different mismatch negativities. *Electroencephalogr. Clin. Neurophysiol.* 78, 466-479.
- Pekkonen, E., Katila, H., Ahveninen, J., Karhu, J., Houttilainen, M., Tiihonen, J. 2002. Impaired temporal lobe processing of preattentive auditory discrimination in schizophrenia. *Schizophr. Bull.* 28, 467-474.
- Petrides, M., Pandya, D.N., 2001. Comparative cytoarchitectonic analysis of the human and the macaque ventrolateral prefrontal cortex and cortico-cortical connection patterns in the monkey. *Eur. J. Neurosci.* 16, 291-310.
- Phillips, C., Rugg, M.D., Friston, K.J., 2002. Systematic regularization of linear inverse solutions of the EEG source localization problem. *Neuroimage* 17, 287-301.
- Picton, T.W., Lins, O.G., Scherg, M., 1995. The recording and analysis of event-related potentials. In: Boller F, Grafman J, editors, *Handbook of Neuropsychology*, Amsterdam: Elsevier, pp 3-73.
- Rey, M-J., Schulz, P., Costa, C., Dick, P., Tissot, R., 1989. Guidelines for the dosage of neuroleptics. 1: Chlorpromazine equivalents of orally administered neuroleptics. *Int Clin Psychopharmacology* 4, 95-104.
- Salisbury, D.F., Farrell, D., Shenton, M.E., Fischer, I.A., Zarate, C., McCarley, R.W. 1999. Mismatch negativity is reduced in chronic but not first episode schizophrenia. *Biol. Psychiatry* 45, 25-26.

- Salisbury, D.F., Shenton, M.E., Griggs, C.B., Bonner-Jackson, A., McCarley, R.W., 2002. Mismatch negativity in chronic schizophrenia and first-episode schizophrenia. *Arch. Gen. Psychiatry* 59, 686-694.
- Sato, Y., Yabe, H., Todd, J., Michie, P.T., Shinozaki, N., Sutoh, T. et al., 2003. Impairment in activation of a frontal attention-switch mechanism in schizophrenic patients. *Biol. Psychol* 62, 49-63.
- Sauer, H., Rosburg, T., Kreitschmann-Andermahr, I., Meier, T., Volz, H-P., Huonker, R. et al., 1997. Impaired mismatch generation in schizophrenics revealed by means of magnetencephalography. *Biol. Psychiatry* 42, 187.
- Scherg, M., Berg, P., 1991. Use of a priori knowledge in brain electromagnetic brain activity. *Brain Topogr.* 4, 143-150.
- Scherg, M., Picton, T.W., 1991. Separation and identification of event-related brain potential components by brain electric source analysis. *Electroencephalogr. Clin. Neurophysiol. Suppl.* 42, 24-37.
- Schröger, E., 1996. A neural mechanism for involuntary attention shifts to changes in auditory stimulation. *J Cogn. Neurosci.* 8, 527-539.
- Shibata, H., Yukie, M., 2003. Differential thalamic connections of the postero-ventral and dorsal posterior cingulate gyrus in the monkey. *Eur. J. Neurosci.* 18, 1615-1626.
- Shinozaki, N., Yabe, H., Sato, Y., Hiruma, T., Sutoh, T., Nashida, T. et al., 2002. The difference in mismatch negativity between the acute and post-acute phase of schizophrenia. *Biol. Psychol.* 59, 105-119.
- Talairach, J., Tournoux, P., 1988. *Co-planar Stereotaxic Atlas of the Human Brain*. New York: Thieme.
- Thompson, P. M., Vidal, C., Giedd, J. N., Cochran, P., Blumenthal J., Nicolson, R. et al., 2001. Mapping adolescent brain change reveals dynamic wave of accelerated grey matter loss in very early-onset schizophrenia. *Proc. Natl. Acad. Sci. (USA)* 98, 11650-11655.
- Tillmann, B., Janata, P., Bharucha, J.J., 2003. Activation of the inferior frontal cortex in musical priming. *Cogn. Brain Res.* 16, 145-161.
- Todd, J., Michie, P.T., Jablensky, A.V., 2002. Do loudness cues contribute to duration mismatch negativity reduction in schizophrenia? *NeuroReport* 12, 4069-4073.
- Umbricht, D., Javitt, D.C., Bates, J., Kane, J., Lieberman, J.A., 2002. Auditory event-related potentials (ERP): indices of both premorbid and illness-related progressive neuropathology in schizophrenia? *Schizophr. Res.* 53, suppl:18.
- Umbricht, D., Koller, R., Schmid, L., Skrabo, A., Grübel, C., Huber, T. et al., 2003. How specific are deficits in mismatch negativity generation to schizophrenia? *Biol. Psychiatry* 53, 1120-1131.
- Valkonen-Korhonen, M., Purhonen, M., Tarkka, I.M., Sipilä, P., Partanen, J., Karhu, J. et al., 2003. Altered auditory consciousness in acutely psychotic, never-medicated first-episode patients. *Cogn. Brain Res.* 17, 747-58.
- Youn, T., Park, H.J., Kim, J-J., Kim, M.S., Kwon, J.S., 2003. Altered hemispheric asymmetry and positive symptoms in schizophrenia: equivalent current dipole of auditory mismatch negativity. *Schizophr. Res.* 59, 253-260.
- Yvert, B., Bertrand, O., Thevenet, M., Echallier, J.F., Pernier, J.A., 1997. Systematic evaluation of the spherical model accuracy in EEG dipole localization. *Electroencephalogr. Clin. Neurophysiol.* 102, 452-459.