



OPEN

PaCO₂-management in the neuro-critical care of patients with subarachnoid hemorrhage

Marvin Darkwah Oppong^{1✉}, Karsten H. Wrede¹, Daniela Müller¹, Alejandro N. Santos¹, Laurèl Rauschenbach¹, Thiemo F. Dinger¹, Yahya Ahmadipour¹, Daniela Pierscianek¹, Mehdi Chihi¹, Yan Li², Cornelius Deuschl², Ulrich Sure¹ & Ramazan Jabbarli¹

The partial pressure of carbon dioxide (PaCO₂) in the arterial blood is a strong vasomodulator affecting cerebral blood flow and the risk of cerebral edema and ischemia after acute brain injury. In turn, both complications are related to poor outcome in patients with aneurysmal subarachnoid hemorrhage (aSAH). We aimed to analyze the effect of PaCO₂ levels on the course and outcome of aSAH. All patients of a single institution treated for aSAH over 13.5 years were included (n = 633). Daily PaCO₂ values from arterial blood gas measurements were recorded for up to 2 weeks after ictus. The study endpoints were: delayed cerebral ischemia (DCI), need for decompressive craniectomy due to increased intracranial pressure > 20 mmHg refractory to conservative treatment and poor outcome at 6-months follow-up (modified Rankin scale > 2). By correlations with the study endpoints, clinically relevant cutoffs for the 14-days mean values for the lowest and highest daily PaCO₂ levels were defined by receiver operating characteristic curve analysis. Association with the study endpoints for the identified subgroups was analyzed using multivariate analysis. The optimal range for PaCO₂ values was identified between 30 and 38 mmHg. ASAH patients with poor initial condition (WFNS 4/5) were less likely to show PaCO₂ values within the range of 30–38 mmHg ($p < 0.001$, OR = 0.44). In the multivariate analysis, PaCO₂ values between 30 and 38 mmHg were associated with a lower risk for decompressive craniectomy ($p = 0.042$, aOR = 0.27), DCI occurrence ($p = 0.035$; aOR = 0.50), and poor patient outcome ($p = 0.004$; aOR = 0.42). The data from this study shows an independent positive association between low normal mean PaCO₂ values during the acute phase of aSAH and patients' outcome. This effect might be attributed to the reduction of intracranial hypertension and alterations in the cerebral blood flow.

The partial pressure of carbon dioxide (PaCO₂) in the arterial blood is a strong vasomodulator affecting the cerebral blood flow (CBF) and subsequently the intracranial pressure (ICP)¹. This cerebrovascular reactivity seems not to be impaired in the case of aneurysmal subarachnoid hemorrhage (aSAH)^{2,3}. Therefore, PaCO₂ alterations might impact the risk of common complications following aneurysmal subarachnoid hemorrhage (aSAH), e.g. delayed cerebral ischemia (DCI) or elevated ICP. Both complications are related to poor outcome in patients with aSAH^{4,5}.

Extreme hypocapnia (PaCO₂ < 30 mmHg) has been historically used in patients with traumatic brain injury (TBI) to lower ICP but is currently abandoned due to its negative impact on outcome⁶. Current guidelines for neuro-critical care of aSAH patients include the management of cerebral vasospasm, blood pressure and increased ICP^{7,8}. However, there are no recommendations on PaCO₂ management and available literature is limited to some small studies. In particular, one surveillance study consisting of 6 patients revealed increased CBF and tissue oxygenation in aSAH patients with increased PaCO₂ levels⁹ while Solaiman and colleagues suggested an association between prolonged hypocapnia and poor patient outcome¹⁰. Moreover, another trial reported on improved outcome after slightly elevated PaCO₂ values during the first 24 h after ictus¹¹.

In the present study, we aimed to analyze the effect of PaCO₂ levels on the course and outcome of aSAH in a large institutional consecutive series.

¹Department of Neurosurgery and Spine Surgery, University Hospital Essen, University of Duisburg-Essen, 45147 Essen, Germany. ²Institute for Diagnostic and Interventional Radiology and Neuroradiology, University Hospital Essen, University of Duisburg-Essen, Essen, Germany. ✉email: marvin.darkwahoppo@uk-essen.de

Methods

Patient population. This is a retrospective study based on the institutional observational aSAH database (registered in the German clinical trial register, Unique identifier: DRKS00008749), which contains all consecutive patients with ruptured intracranial aneurysms treated at our institution between 01/2003 and 06/2016. The exclusion criteria were: (a) treatment due to a mycotic aneurysm; (b) admission > 48 h after ictus; (c) no treatment of the ruptured aneurysm; (d) hospital stay duration < 72 h after admission.

The approval of the institutional ethics committee (Ethik-Kommission, Medizinische Fakultät der Universität Duisburg-Essen, Registration number: 15-6331-BO) for this study was obtained and has been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments. The study was registered in the German clinical trial registry (DRKS, Unique identifier: DRKS00008749). All patients or their relatives gave written consent within the treatment contract before inclusion into the database.

SAH management. All patients were initially admitted at our neurosurgical intensive care unit for at least 10 days after ictus followed by further surveillance at our intermediate care unit for at least 4 days, unless the cases with shorter survival. Acute hydrocephalus was treated by placement of an external ventricular drainage (EVD). Further conservative aSAH management was performed according to the latest guidelines^{7,8} and included oral nimodipine and euvolemia maintenance for 3 weeks. Conservative and invasive treatment of cerebral vasospasm after aSAH has been described in detail elsewhere¹². The Bleeding source was confirmed using digital subtraction and/or computed tomography (CT)-angiography. Treatment decision was made upon interdisciplinary discussion between the interventional neuroradiologist and the vascular neurosurgeon on call. Monitoring included daily transcranial Doppler ultrasound, ICP monitoring via EVD and 3 hourly blood gas samples taken from the arterial line. Mechanical ventilation (MV) was induced for aneurysm treatment and discontinued as soon as possible depending on the patient's pulmonary and clinical state.

All patients underwent a routine postoperative CT scan within 24 h after aneurysm treatment. Moreover, additional CT imaging was performed in cases of any neurological deterioration or prolonged impairment in the conscious state.

Conservative and surgical ICP management. In case of sustained ICP increase > 20 mmHg, conservative treatment was initiated maintaining cerebral perfusion pressure at > 60 mmHg as follows:

1. Level 1: Infusion and catecholamines, normothermia, elevation of the head to 30°, deep sedation, osmotherapy with 20% mannitol and forced cerebrospinal fluid drainage.
2. Level 2: Relaxation, application of tromethamine, and barbiturate coma, guided by a burst-suppression-EEG pattern.

In case of persisting ICP elevation refractory to conservative treatment, decompressive craniectomy (DC) was performed. There was no specific target area for PaCO₂ levels regarding ICP management.

Data management. The digital archive and imaging were reviewed for demographic, clinical, laboratory and all radiographic parameters. To assess the value of PaCO₂ for the course and outcome of aSAH, daily highest and lowest PaCO₂ values during the first 14 days after ictus were recorded.

The clinical severity of aSAH was documented using the world federation of neurological surgery grading system (WFNS) scale¹³. For statistical analysis, we dichotomized patients into good (WFNS 1–3) and poor grade (WFNS grade 4–5) cases. For assessment of radiographic severity, the original Fisher scale was applied¹⁴, with further dichotomization into low (Fisher scale 1 and 2) and high (Fisher scale 3 and 4) grades.

Infarctions that were not related to surgical approach or intracerebral hemorrhage were documented as early infarction if present ≤ 48 h after early aneurysm occlusion and as delayed cerebral ischemia (DCI) if occurring later than 48 h after aneurysm occlusion¹⁵. All imaging data was reviewed by the last author (RJ) blinded at that time for any clinical information.

The total time of MV in days was documented. A MV > 7 days was defined as prolonged MV. Pneumonia was diagnosed by chest x-ray in case of pulmonary alterations and/or elevated laboratory infectious parameters. Treatment was performed according to the German guidelines for nosocomial pneumonia¹⁶.

The functional outcome was measured upon the routine outpatient follow-ups performed at 6 months after aSAH using the modified Rankin scale (mRS)¹⁷. A mRS greater than 2 was defined as poor outcome.

Statistical analysis. The primary study endpoints were the association between PaCO₂ values and: (1) the risk of poor outcome at 6 months after aSAH, (2) occurrence of DCI, and (3) the need for DC refractory to conservative treatment. PaCO₂ values were additionally correlated with the patients' baseline characteristics. SPSS 22 for Windows (IBM Corp.) was used for all statistical analysis. The significance level was set to $p < 0.05$.

According to the correlations with the primary study endpoints, clinically relevant cutoffs for the 14-days mean values for the lowest and highest daily PaCO₂ levels were defined by ROC analysis. After that the whole cohort was divided into two groups:

1. Patients with PaCO₂ values within the range of the ROC-defined cutoffs were referred to the “*optimal PaCO₂*”-group;
2. The remaining aSAH patients with records exceeding the PaCO₂ limits (lower than the lowest and/or higher than the highest cutoff) were referred to the “*non-optimal PaCO₂*”-group.




Figure 1. Flow-Chart of recruitment process for this study.

Endpoint	Mean lowest PaCO ₂ (mmHg)		Mean highest PaCO ₂ (mmHg)	
Poor outcome (yes/no)	31.57 ± 4.55	32.04 ± 4.79	40.20 ± 3.75	38.85 ± 4.09
DCI (yes/no)	31.16 ± 3.84	32.00 ± 4.89	40.02 ± 3.70	39.33 ± 4.06
DC due to refractory ICP increase (yes/no)	30.84 ± 2.87	32.04 ± 5.27	40.12 ± 3.56	39.22 ± 4.03

Table 1. Overview of lowest/highest mean PaCO₂ value distribution among the different endpoints.

The associations between the allocation to the PaCO₂-related groups and the study endpoints were assessed in a multivariable manner using binary logistic regression analysis for the study endpoints adjusted for endpoint-relevant confounders. To analyze of the need for DC in cases refractory to conservative ICP treatment, the cases with DC performed at admission prior to the initiation of conservative ICP treatment (n = 145) were excluded. Missing data was managed utilizing multiple imputations.

Results

Patient population. A total of 633 patients were eligible for this study (see Fig. 1). The majority of patients were females (n = 422; 66.7%) and presented with severe aSAH on the initial CT scan (n = 538; 91.2%). About half of the patients were at WFNS grade 4–5 at admission (n = 300; 47.4%). DCI occurred in one of four patients (n = 153; 24.2%) and 49.5% of all cases (n = 311) revealed poor outcome at 6-month follow-up. See also Table 1 in the supplemental material for a detailed overview of the cohort characteristics.

Identification of clinically relevant lowest/highest PaCO₂ cutoffs and allocation of the patients to the PaCO₂-related groups. In accordance with the ROC analysis investigating the 14-days lowest/highest mean PaCO₂ levels and the study endpoints, the clinically relevant cutoff for the lowest 14-days mean PaCO₂ was set at 30 mmHg, and the cutoff for the highest 14-days mean PaCO₂ at 38 mmHg. Accordingly, aSAH patients with PaCO₂ values within the above-mentioned ranges (n = 109) were referred to the “*optimal PaCO₂*” group, whereas the remaining individuals (n = 524) were included in the “*non-optimal PaCO₂*” group. See also Figs. 1, 2 and Table 2 in the supplemental material. A detailed listing of mean PaCO₂ values for the different endpoints is given in Table 1.

Parameter	Optimal PaCO ₂	Non optimal PaCO ₂	<i>p</i>	OR	95% CI
	%/mean ± SD	%/mean ± SD			
Age (years)	52.7 ± 14.0	54.9 ± 13.6	0.136	–	–
Sex female	67.9%	66.4%	0.766	1.07	0.69–1.66
WFNS 4/5	31.2%	50.8%	< 0.001	0.44	0.28–0.68
Fisher 3/4	90.4%	91.3%	0.777	0.90	0.42–1.91
Treatment clipping	37.6%	42.0%	0.399	0.83	0.55–1.27

Table 2. Statistical analysis of demographic and clinical baseline characteristics for optimal and non-optimal PaCO₂ groups. Statistically significant *p*-values are marked bold.

Parameter	<i>p</i>	aOR	95% CI
WFNS 4/5	0.158	1.64	0.83–3.27
Fisher 3/4	0.715	1.31	0.31–5.50
Age (> 65 years)	0.017	0.30	0.11–0.81
Conservative ICP therapy level 1	< 0.001	41.3	9.79–174.23
Conservative ICP therapy level 2	0.357	1.72	0.54–5.45
Optimal PaCo ₂ (30–38 mmHg)	0.042	0.27	0.08–0.96

Table 3. Multivariate analysis of factors affecting the need for DC. Statistically significant *p*-values are marked bold.




Figure 2. Distribution of risk for (A) DC due to refractory ICP increase, (B) DCI and (C) poor outcome between the optimal PaCO₂ and the non-optimal PaCO₂ group.

PaCO₂-groups and baseline characteristics. ASAH individuals with poor initial clinical condition (WFNS grade 4–5) were less likely to show PaCO₂ values within the optimal range ($p < 0.001$; odds ratio [OR] = 0.44; 95% confidence interval [95%CI] 0.28–0.68). Other baseline demographic and clinical characteristics of the patients showed no significant associations with the PaCO₂-groups (Table 2).

Surgical ICP management. Individuals in the “optimal PaCO₂” group showed less frequently an ICP increase unresponsive to conservative therapy. Accordingly, this patient group was less likely to require DC ($p = 0.042$; adjusted [a]OR = 0.27; 95%CI 0.08–0.96, see Table 3 and Fig. 2A).

DCI and vasospasm risk. The DCI’s overall occurrence rate was significantly reduced in the “optimal PaCO₂” group (Fig. 2B). This effect of PaCO₂ on DCI occurrence was independent of clinical/radiographic aSAH severity, age, vasospasm, pneumonia and prolonged MV ($p = 0.035$; aOR = 0.50; 95%CI 0.26–0.95; Table 4).

Parameter	<i>p</i>	aOR	95% CI
WFNS 4/5	0.007	1.80	1.17–2.77
Fisher 3/4	0.112	2.51	0.08–7.86
Age (> 65 years)	0.104	1.46	0.92–2.32
Vasospasm	< 0.001	2.89	1.97–4.46
Pneumonia	0.006	1.87	1.20–2.93
Prolonged MV (> 7 days)	0.417	1.20	0.77–1.86
Optimal PaCO ₂ (30–38 mmHg)	0.035	0.50	0.26–0.95

Table 4. Multivariate analysis of factors impacting the occurrence of DCI and poor outcome. Statistically significant *p*-values are marked bold.

Parameter	<i>p</i>	aOR	95% CI
WFNS 4/5	< 0.001	3.27	2.14–5.00
Fisher 3/4	0.011	3.61	1.35–9.65
Age (> 65 years)	< 0.001	3.30	2.00–5.45
DCI	< 0.001	6.45	3.78–11.00
Pneumonia	0.006	2.10	1.24–3.57
Prolonged MV (> 7 days)	> 0.001	2.78	1.83–4.23
Optimal PaCO ₂ (30–38 mmHg)	0.004	0.42	0.23–0.76

Table 5. Multivariate analysis of factors impacting the occurrence of poor outcome. Statistically significant *p*-values are marked bold.

Functional outcome. Patients in the “optimal PaCO₂” group showed significantly reduced risk for a poor outcome at 6-month follow-up in multivariate analysis: *p* = 0.004; aOR = 0.42; 95%CI 0.23–0.76 (see Table 5 and Fig. 2C).

Discussion

This study aimed to evaluate the influence of PaCO₂ levels on major complications (like occurrence of DCI and intractable ICP increase) and functional outcome of aSAH. Therefore, we defined an optimal PaCO₂ window between moderate hypocapnia (≥ 30 mmHg) and slightly elevated normocapnia (≤ 38 mmHg). aSAH individuals with PaCO₂ values during the first 2 weeks within this range were at lower risk of DCI, DC due to conservatively intractable ICP increase and poor outcome 6 months after aSAH.

PaCO₂ and ICP elevations refractory to conservative therapy. In case of malignant infarction, DC is a common neurosurgical treatment option for sustained ICP increase refractory to conservative management¹⁸. In contrast to stroke, no clear guidelines regarding the indication and timing of DC exist in the setting of aSAH¹⁹. Patients that develop sustained ICP increase necessitating DC often face a poor outcome^{19–21}. Therefore, early prevention of such intractable ICP increase is of paramount importance for aSAH patients’ outcome.

It is generally acknowledged that the decreasing of PaCO₂ leads to lower ICP values²² and the pathophysiological background of this effect is related to arteriolar vasoconstriction, especially in the pial and cortical vessels. This mechanism reduces CBF and is believed to subsequently reduce ICP by reducing the development of cerebral edema^{23,24}. Historically, extreme hypocapnia (< 30 mmHg) was commonly used to lower ICP in patients with TBI⁶. However, this practice was later abandoned due to poor functional outcome of individuals treated with PaCO₂ < 30 mmHg⁶. Of note, the ICP-lowering effect of PaCO₂ might be present also at moderate hypocapnia²³. In our cohort, individuals with consistently low-normal PaCO₂ values ranging between 30 and 38 mmHg were at lower risk for ICP increase unresponsive to conservative therapy than the remaining aSAH cohort. Therefore, this PaCO₂ management might become a useful component of conservative ICP treatment in aSAH.

PaCO₂ and DCI risk. The occurrence of DCI has a paramount impact on the course and outcome of aSAH²⁵. DCI’s exact pathophysiology in aSAH has not yet been completely understood, but multi-factorial genesis of DCI is supposed²⁶. Cerebral vasospasm is one major mechanism of DCI and aSAH management includes nimodipine application, induced hypertension and angiographic vasospasmolysis^{25,27}. Despite cerebral vasospasm, inflammatory factors, micro-thrombosis and cortical spreading depolarization seem to play a role in DCI development^{4,28–31}.

Strong alterations of PaCO₂ in both directions might also contribute to DCI risk after aSAH. On one side, hypercapnia is associated with ICP increase and subsequent decrease of cerebral perfusion pressure and CBF. On the other side, hypocapnia also results in CBF decline due to vasoconstriction. At the same time, induced

hypertension initiated in patients with cerebral vasospasm might alleviate the vasoconstrictory effect of hypocapnia on CBF. Therefore, the negative clinical impact of hypocapnia observed in TBI cohorts is not to be expected to the same extent in aSAH. Instead, moderate hypocapnia and induced hypertension potentially present an optimal framework condition for the prevention of DCI. This constellation might explain our results showing that aSAH patients with PaCO₂ values ranging between 30 and 38 mmHg were at lower DCI risk. This independent association between PaCO₂ and DCI risk encourages further research of the potential link between PaCO₂, CBF and DCI in aSAH patients.

PaCO₂ and outcome following aSAH. Our data underlines that moderately decreased PaCO₂ independently reduces the risk of poor outcome following aSAH. In TBI and cerebrovascular accidents, aggressive hypocapnia has been associated with poor outcome and mortality^{32–34}. At least for TBI, the critical cut-off was reported at the values between 27 and 32 mmHg^{33,34}. Also no relevant advantage could be shown for extreme hypocapnia (30 mmHg) in TBI treatment⁶. Comparable results exist for stroke³⁵. Interestingly, in a subgroup analysis excluding major vessel occlusion and mainly including hemorrhagic stroke, hypocapnia showed a tendency to lower mortality and better outcome³⁵. Data from aSAH are very sparse and suggest poorer outcome in patients regularly having PaCO₂ levels below 35 mmHg¹⁰. This discrepancy to our results might be related to the fact that the “< 35 mmHg PaCO₂” group in this study¹⁰ also included individuals with PaCO₂ values < 30 mmHg. Recent published data suggests that PaCO₂ values regularly reaching high normal values (> 40 mmHg) are harmful for aSAH patients³⁶.

In summary, the positive effect of “low-normal PaCO₂” management during the first 2 weeks after aneurysm rupture on the risk of DCI, intractable ICP increase and poor outcome after aSAH might speak for the implementation of specific PaCO₂ target area in future guidelines for the management of aSAH patients requiring MV, provided the confirmation of our results in a prospective controlled trial.

Limitations

This study's main drawback is its retrospective design, resulting in lower accuracy and consistency of data than in comparable prospective collected data sets. This disadvantage might be partly outweighed by the cohort's size, strict exclusion criteria and robust statistical assessment. Our data does not allow a direct correlation between PaCO₂ level and corresponding ICP. This connection was only made indirectly by using the clinical surrogate parameters of increased ICP, i.e. conservative and surgical ICP therapy. Furthermore, this study does not exclusively include patients that were ventilated for the whole surveillance period. To address this potential risk of bias, the results were corrected for the duration of ventilation.

Conclusions

Our data suggests that patients with aSAH might profit from low-normal PaCO₂ values during the first weeks after aneurysm rupture. This effect is highlighted by reducing DCI and intracranial hypertension, resulting in lower rate of poor outcome in aSAH individuals with PaCO₂ values ranging between 30 and 38 mmHg in our cohort. The results from this study urge for a prospective evaluation of the clinical benefit of low-normal PaCO₂ values in the treatment of aSAH.

Received: 5 May 2021; Accepted: 6 September 2021

Published online: 28 September 2021

References

- Ainslie, P. N. & Duffin, J. Integration of cerebrovascular CO₂ reactivity and chemoreflex control of breathing: Mechanisms of regulation, measurement, and interpretation. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **296**, R1473–1495. <https://doi.org/10.1152/ajpregu.91008.2008> (2009).
- Romner, B. *et al.* Simultaneous transcranial Doppler sonography and cerebral blood flow measurements of cerebrovascular CO₂-reactivity in patients with aneurysmal subarachnoid haemorrhage. *Br. J. Neurosurg.* **5**, 31–37. <https://doi.org/10.3109/02688699108998444> (1991).
- Diringer, M. N., Kirsch, J. R., Hanley, D. F. & Traystman, R. J. Altered cerebrovascular CO₂ reactivity following subarachnoid hemorrhage in cats. *J. Neurosurg.* **78**, 915–921. <https://doi.org/10.3171/jns.1993.78.6.0915> (1993).
- DarkwahOppong, M. *et al.* Post-treatment antiplatelet therapy reduces risk for delayed cerebral ischemia due to aneurysmal subarachnoid hemorrhage. *Neurosurgery* <https://doi.org/10.1093/neuros/nyy550> (2018).
- Mohme, M. *et al.* Irreversible total loss of brain function and organ donation in patients with aneurysmal subarachnoid hemorrhage. *World Neurosurg.* <https://doi.org/10.1016/j.wneu.2017.06.024> (2017).
- Muizelaar, J. P. *et al.* Adverse effects of prolonged hyperventilation in patients with severe head injury: A randomized clinical trial. *J. Neurosurg.* **75**, 731–739. <https://doi.org/10.3171/jns.1991.75.5.0731> (1991).
- Connolly, E. S. Jr. *et al.* Guidelines for the management of aneurysmal subarachnoid hemorrhage: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke; J. Cereb. Circ.* **43**, 1711–1737. <https://doi.org/10.1161/STR.0b013e3182587839> (2012).
- Steiner, T. *et al.* European stroke organization guidelines for the management of intracranial aneurysms and subarachnoid haemorrhage. *Cerebrovasc. Dis. (Basel, Switz.)* **35**, 93–112. <https://doi.org/10.1159/000346087> (2013).
- Westermaier, T. *et al.* Controlled transient hypercapnia: A novel approach for the treatment of delayed cerebral ischemia after subarachnoid hemorrhage?. *J. Neurosurg.* **121**, 1056–1062. <https://doi.org/10.3171/2014.7.jns132611> (2014).
- Solaiman, O. & Singh, J. M. Hypocapnia in aneurysmal subarachnoid hemorrhage: Incidence and association with poor clinical outcomes. *J. Neurosurg. Anesthesiol.* **25**, 254–261. <https://doi.org/10.1097/ANA.0b013e3182806465> (2013).
- Lång, M. *et al.* Early moderate hyperoxemia does not predict outcome after aneurysmal subarachnoid hemorrhage. *Neurosurgery* **78**, 540–545. <https://doi.org/10.1227/neu.000000000001111> (2016).
- Jabbarli, R. *et al.* Endovascular treatment of cerebral vasospasm after subarachnoid hemorrhage: More is more. *Neurology* <https://doi.org/10.1212/wnl.0000000000007862> (2019).

13. Teasdale, G. M. *et al.* A universal subarachnoid hemorrhage scale: Report of a committee of the world federation of neurosurgical societies. *J. Neurol. Neurosurg. Psychiatr.* **51**, 1457 (1988).
14. Fisher, C. M., Kistler, J. P. & Davis, J. M. Relation of cerebral vasospasm to subarachnoid hemorrhage visualized by computerized tomographic scanning. *Neurosurgery* **6**, 1–9 (1980).
15. Vergouwen, M. D. *et al.* Definition of delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage as an outcome event in clinical trials and observational studies: Proposal of a multidisciplinary research group. *Stroke; J. Cereb. Circ.* **41**, 2391–2395. <https://doi.org/10.1161/strokeaha.110.589275> (2010).
16. Lorenz, J. *et al.* Nosocomial pneumonia: Prevention, diagnosis, treatment. *Pneumol. (Stuttg., Ger.)* **57**, 532–545. <https://doi.org/10.1055/s-2003-42217> (2003).
17. van Swieten, J. C., Koudstaal, P. J., Visser, M. C., Schouten, H. J. & van Gijn, J. Interobserver agreement for the assessment of handicap in stroke patients. *Stroke; J. Cereb. Circ.* **19**, 604–607 (1988).
18. Warner, J. J., Harrington, R. A., Sacco, R. L. & Elkind, M. S. V. Guidelines for the early management of patients with acute ischemic stroke: 2019 update to the 2018 guidelines for the early management of acute ischemic stroke. *Stroke; J. Cereb. Circ.* **50**, 3331–3332. <https://doi.org/10.1161/strokeaha.119.027708> (2019).
19. DarkwahOppong, M. *et al.* Decompressive craniectomy in aneurysmal subarachnoid hemorrhage: Who and when?—A systematic review and meta-analysis. *Clin. Neurol. Neurosurg.* **199**, 106252. <https://doi.org/10.1016/j.clineuro.2020.106252> (2020).
20. Jabbarli, R. *et al.* Time is brain! Analysis of 245 cases with decompressive craniectomy due to subarachnoid hemorrhage. *World Neurosurg.* **98**, 689–694 e682. <https://doi.org/10.1016/j.wneu.2016.12.012> (2017).
21. Jabbarli, R. *et al.* The pressure score to predict decompressive craniectomy after aneurysmal subarachnoid haemorrhage. *Brain Commun.* **2**, fcaa134. <https://doi.org/10.1093/braincomms/fcaa134> (2020).
22. Raichle, M. E. & Plum, F. Hyperventilation and cerebral blood flow. *Stroke; J. Cereb. Circ.* **3**, 566–575. <https://doi.org/10.1161/01.str.3.5.566> (1972).
23. Sadoshima, S. *et al.* Response of cortical and pial arteries to changes of arterial CO₂ tension in rats—A morphometric study. *Brain Res.* **189**, 115–120. [https://doi.org/10.1016/0006-8993\(80\)90011-6](https://doi.org/10.1016/0006-8993(80)90011-6) (1980).
24. Tuor, U. I. & Farrar, J. K. Pial vessel caliber and cerebral blood flow during hemorrhage and hypercapnia in the rabbit. *Am. J. Physiol.* **247**, H40–H51. <https://doi.org/10.1152/ajpheart.1984.247.1.H40> (1984).
25. Francoeur, C. L. & Mayer, S. A. Management of delayed cerebral ischemia after subarachnoid hemorrhage. *Crit. Care (Lond., Engl.)* **20**, 277. <https://doi.org/10.1186/s13054-016-1447-6> (2016).
26. Geraghty, J. R. & Testai, F. D. Delayed cerebral ischemia after subarachnoid hemorrhage: Beyond vasospasm and towards a multifactorial pathophysiology. *Curr. Atheroscler. Rep.* **19**, 50. <https://doi.org/10.1007/s11883-017-0690-x> (2017).
27. Etminan, N., Vergouwen, M. D., Ilodigwe, D. & Macdonald, R. L. Effect of pharmaceutical treatment on vasospasm, delayed cerebral ischemia, and clinical outcome in patients with aneurysmal subarachnoid hemorrhage: A systematic review and meta-analysis. *J. Cereb. Blood Flow Metabol.: Off. J. Int. Soc. Cereb. Blood Flow Metabol.* **31**, 1443–1451. <https://doi.org/10.1038/jcbfm.2011.7> (2011).
28. Frontera, J. A. *et al.* The role of platelet activation and inflammation in early brain injury following subarachnoid hemorrhage. *Neurocrit. Care* **26**, 48–57. <https://doi.org/10.1007/s12028-016-0292-4> (2017).
29. McBride, D. W., Blackburn, S. L., Peeyush, K. T., Matsumura, K. & Zhang, J. H. The role of thromboinflammation in delayed cerebral ischemia after subarachnoid hemorrhage. *Front. Neurol.* **8**, 555. <https://doi.org/10.3389/fneur.2017.00555> (2017).
30. Foreman, B. The pathophysiology of delayed cerebral ischemia. *J. Clin. Neurophysiol.: Off. Publ. Am. Electroencephalogr. Soc.* **33**, 174–182. <https://doi.org/10.1097/wnp.0000000000000273> (2016).
31. Jabbarli, R. *et al.* Laboratory biomarkers of delayed cerebral ischemia after subarachnoid hemorrhage: A systematic review. *Neurosurg. Rev.* <https://doi.org/10.1007/s10143-018-1037-y> (2018).
32. Rout, M. W., Lane, D. J. & Wollner, L. Prognosis in acute cerebrovascular accidents in relation to respiratory pattern and blood gas tensions. *BMJ* **3**, 7–9. <https://doi.org/10.1136/bmj.3.5765.7> (1971).
33. Davis, D. P. *et al.* The impact of hypoxia and hyperventilation on outcome after paramedic rapid sequence intubation of severely head-injured patients. *J. Trauma* **57**, 1–8; discussion 8–10. <https://doi.org/10.1097/01.ta.0000135503.71684.c8> (2004).
34. Davis, D. P. *et al.* Early ventilation and outcome in patients with moderate to severe traumatic brain injury. *Crit. Care Med.* **34**, 1202–1208. <https://doi.org/10.1097/01.ccm.0000208359.74623.1c> (2006).
35. Christensen, M. S. *et al.* Cerebral apoplexy (stroke) treated with or without prolonged artificial hyperventilation. 1. Cerebral circulation, clinical course, and cause of death. *Stroke; J. Cereb. Circ.* **4**, 568–631. <https://doi.org/10.1161/01.str.4.4.568> (1973).
36. Reiff, T., Barthel, O., Schönenberger, S. & Mundiyanapurath, S. High-normal P(a)CO₂ values might be associated with worse outcome in patients with subarachnoid hemorrhage—A retrospective cohort study. *BMC Neurol.* **20**, 31. <https://doi.org/10.1186/s12883-020-1603-0> (2020).

Author contributions

All authors contributed to the study conception and design. Material preparation and analysis were performed by M.D.O. and R.J., Fig. 2 was prepared by L.R. Data collection was performed by M.D.O. and R.J. The first draft of the manuscript was written by M.D.O. and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-98462-2>.

Correspondence and requests for materials should be addressed to M.D.O.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021

DuEPublico

Duisburg-Essen Publications online

UNIVERSITÄT
DUISBURG
ESSEN

Offen im Denken

ub | universitäts
bibliothek

This text is made available via DuEPublico, the institutional repository of the University of Duisburg-Essen. This version may eventually differ from another version distributed by a commercial publisher.

DOI: 10.1038/s41598-021-98462-2

URN: urn:nbn:de:hbz:465-20240905-113244-8



This work may be used under a Creative Commons Attribution 4.0 License (CC BY 4.0).