



OPEN ACCESS

EDITED BY

Helmi Chaabene,
Otto von Guericke University Magdeburg,
Germany

REVIEWED BY

Arzu Genc,
Dokuz Eylul University, Türkiye
Dario Martínez-García,
University of Granada, Spain

*CORRESPONDENCE

Johanna Lambrich
✉ johanna.lambrich@uni-due.de

RECEIVED 18 February 2025

ACCEPTED 11 April 2025

PUBLISHED 29 April 2025

CITATION

Lambrich J and Muehlbauer T (2025) Effects of fatigue on physiological, physical fitness, and stroke performance related parameters in healthy tennis players: a systematic review and meta-analysis.

Front. Sports Act. Living 7:1578914.

doi: 10.3389/fspor.2025.1578914

COPYRIGHT

© 2025 Lambrich and Muehlbauer. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Effects of fatigue on physiological, physical fitness, and stroke performance related parameters in healthy tennis players: a systematic review and meta-analysis

Johanna Lambrich* and Thomas Muehlbauer

Division of Movement and Training Sciences/Biomechanics of Sport, University of Duisburg-Essen, Essen, Germany

Fatigue is a multifactorial phenomenon involving central and peripheral mechanisms that could negatively affect performance-related measures in tennis players. The aim of this systematic review and meta-analysis was to quantify the effects of fatigue on physiological (e.g., blood lactate, heart rate), physical fitness (e.g., explosive muscle strength, speed), and stroke performance (e.g., stroke velocity or accuracy) related parameters in healthy tennis players and to provide insight into training and recovery strategies. A systematic literature search of PubMed, Web of Science, and SportDiscus identified studies that examined effects of fatigue in tennis. Inclusion criteria required that studies were conducted with healthy participants, applied fatigue-inducing interventions, and used pre-/post-test designs measuring physiological, physical, or stroke performance related parameters. Effect sizes were calculated using weighted standardized mean differences (*SMD*) to assess the impact of fatigue. The search identified a total of $N = 642$ records. Eighteen trials (318 tennis players) were included. Fatigue evoked large to moderate negative effect on physiological ($SMD_w = -4.19$), physical fitness ($SMD_w = -0.74$), and stroke performance ($SMD_w = -0.60$) related parameters. The larger negative effects of fatigue on physiological and physical parameters compared to stroke performance-related outcomes indicate the importance of targeted recovery strategies (e.g., hydration, nutrition or cold baths). For stroke performance, non-fatigued states are recommended for learning new skills, while practice under fatigued conditions may help to maintain biomechanical efficiency during prolonged games.

KEYWORDS

racket sport, exhaustion, physiology, physical fitness, stroke performance

Introduction

Fatigue is generally defined as the loss of the ability to maintain a certain level of performance or strength over an extended period (1). It is a multifactorial phenomenon involving both central and peripheral mechanisms. Central fatigue affects neural control in the central nervous system, reducing signal transmission to the muscles. Peripheral fatigue, on the other hand, refers to muscular processes such as the accumulation of metabolic waste products or reduced calcium release that impair muscle contractility

(2, 3). These peripheral impairments directly contribute to motor performance fatigue by reducing the muscle's ability to generate or sustain force during repeated or prolonged efforts. Motor performance fatigue specifically refers to a decline in the neuromuscular system's ability to sustain force production, resulting from both central and peripheral mechanisms. This type of fatigue is critical for understanding the relationship between physiological processes and performance during motor tasks, as it directly affects task performance and endurance (4, 5). Recent evidence (6, 7) suggests that central and peripheral mechanisms do not operate in isolation but interact dynamically during prolonged or high-intensity exercise. For example, peripheral metabolic disturbances may enhance central fatigue through afferent feedback pathways, while central drive may modulate the extent of peripheral muscle activation and fatigue development.

Studies have shown that in tennis players fatigue leads to biomechanical deviations in stroke technique (e.g., ball impact height during serve) and results in a decline in stroke speed and accuracy (8, 9). It has also been shown that fatigue reduces footwork and trunk stability, which in turn impairs the efficiency of the stroke technique (10). At the same time, changes in muscle activation and kinematics can lead to an increased susceptibility to injury, especially during repetitive, high-speed movements such as serves (11, 12). Fatigue has also been shown to increase the error rate of serves and defensive shots (13–15). For example, it has been shown that stroke accuracy decreases by up to 49.6% under high-intensity training, even in experienced tennis players (16). These biomechanical and technical impairments may have a direct impact on match results, as reduced stroke speed and accuracy can increase unforced error rates, limit tactical options, and compromise the ability to execute aggressive plays while fatigued (17). In addition, fatigue can lead to wrong tactical decisions (18).

Although there have been several reviews on fatigue in tennis (18–20), no study has simultaneously examined parameters related to physiology, physical fitness, and stroke performance. Existing reviews tend to focus on isolated aspects—such as the physiological mechanisms underlying fatigue or their biomechanical effects on stroke execution—without examining how these components interact under real-world performance conditions. In competitive tennis, however, fatigue is a multifactorial phenomenon that affects multiple dimensions of performance simultaneously. For example, a reduction in physiological efficiency (e.g., elevated lactate levels or reduced cardiovascular output) can compromise neuromuscular function, leading to impaired movement quality and reduced stroke precision. Therefore, a holistic synthesis is essential to understand how fatigue initiates systemic cascades that contribute to performance decline. While previous reviews have relied primarily on narrative summaries or focused on single domains, the present meta-analysis is the first to quantitatively integrate physiological, physical fitness and stroke performance parameters within a unified analytical framework. In addition, by including subgroup comparisons between elite and sub-elite players, this

study provides novel insights into how fatigue responses may vary according to competitive level.

The aim of this systematic review and meta-analysis was to analyze and quantify the effects of motor performance fatigue (4) on physiological parameters, physical performance and stroke performance in healthy tennis players. We hypothesized that performance in all three categories—physiological, physical fitness and stroke performance—would be reduced due to motor fatigue. Furthermore, we expected elite players to show greater resilience to fatigue, particularly in physiological parameters such as blood lactate and heart rate, and physical fitness outcomes such as countermovement jump and sprint performance. The findings of this review may assist coaches and sport scientists in developing more targeted training and recovery strategies that address the specific fatigue-related limitations in physiological, physical fitness and stroke performance parameters. By identifying which areas of performance are most affected by fatigue and how these effects differ between performance levels, practitioners can implement individualized interventions to optimize race readiness, maintain technical execution under pressure and reduce the risk of overuse injuries.

Methods

Search strategy

A systematic literature search of the PubMed, Web of Science, and SportDiscus databases was conducted to identify eligible articles. The following Boolean expression was used:

```
tennis AND (fatigue AND (functional OR performance OR
agility OR flexibility OR athletic OR strength OR power OR
speed OR fitness OR physical OR reaction time OR stroke
OR serve OR forehand OR backhand OR balance OR
resistance OR physiology OR heart rate OR blood lactate OR
creatine OR cardiovascular)) NOT table
```

The search was conducted across the entire history of each database, up to and including January 2025. Only articles written in English with full-text access were included. No search filters regarding publication type, study design, or date were applied in order to maximize the sensitivity and comprehensiveness of the literature search. Furthermore, the reference lists of the included studies and relevant reviews were examined to identify additional eligible articles. After removing duplicates, the titles and abstracts of all retrieved records were evaluated for eligibility based on the inclusion and exclusion criteria independently by both authors (Table 1). The full texts of potentially eligible studies were then assessed independently, with any discrepancies resolved through discussion and consensus. The process of the literature search, study selection, and exclusion of articles is summarized in a PRISMA flow chart (21) (see Figure 1). Disagreements during the study selection process were resolved through discussion, with unresolved cases adjudicated by a third, independent reviewer affiliated with the host institution.

TABLE 1 Overview of the inclusion and exclusion criteria.

Category	Inclusion criteria	Exclusion criteria
Population	Healthy female and male tennis players	Injured tennis players; no tennis players; participants with existing physical or health limitations
Intervention	Studies inducing fatigue, e.g., through repeated sprints, endurance tests, or tennis game simulations; research assessing physiological, physical, or stroke performance of fatigue.	Studies without a motor performance fatigue-related intervention, e.g., cognitive fatigue
Comparison	Pre-/post-test designs	Studies lacking comparative conditions
Outcome	At least one physical fitness, physiological or stroke performance related parameter	Data did not allow calculating effect size
Study design	Experimental studies, cross-sectional studies, or cohort studies, intervention studies with posttest after <24 h	Theoretical work, opinion articles, or studies without primary data or post-testing after more than 24 h

Study selection criteria

The inclusion and exclusion criteria are summarized in Table 1. Studies were eligible for this review if they (a) included healthy female or male tennis players, (b) implemented a motor performance fatigue-inducing intervention such as repeated sprints, endurance tests, or simulated tennis matches, (c) employed a pre-/post-test design, (d) reported at least one parameter of physiological response, physical fitness, or stroke performance, and (e) conducted interventions with a short-term focus (post-test after less than 24 h). Studies were excluded if (a) injured tennis players, non-tennis players, or participants with physical or health limitations were examined, (b) the intervention lacked a defined method for inducing fatigue, focused solely on long-term adaptations, or addressed only mental fatigue, (c) no pre-/post-test design or comparative condition was present, (d) the data did not allow the calculation of effect sizes, or (e) post-test measurements were conducted after more than 24 h. To ensure that the observed effects reflect acute motor fatigue, only studies with a post-test interval of less than 24 h were included. This decision is supported by the findings of Girard et al. (22), who demonstrated that impairments in physical performance induced by match play were no longer detectable after 24 h of recovery.

Assessment of methodological quality


The quality of the included studies was assessed using the Joanna Briggs Institute (JBI) Critical Appraisal Tool for Case Series Studies, as described by Munn et al. (23). The tool consists of ten items, each designed to assess critical aspects of methodological quality. These items are answered with “yes”, “no” or “unclear”. Four questions (1, 2, 3, 4) assess inclusion criteria and clarity of case series reporting, focusing on whether inclusion criteria were clearly defined and appropriately applied. Another three questions (5, 6, 7) address the potential for bias in patient selection, whether consecutive cases were included, and the completeness of the clinical information provided. The remaining three questions (8, 9, 10) address the appropriateness of the statistical methods and the reporting of results. Discrepancies in the quality assessment were resolved by a third party, a graduate sports scientist associated with the host institution.

Additionally, Egger’s test was applied to assess the presence of publication bias, as it provides a regression-based approach to detect asymmetry in funnel plots. Egger’s test is widely used to

detect small study effects and publication bias in meta-analyses, although its reliability decreases in cases of high heterogeneity (24). To assess and adjust for potential publication bias, Duval and Tweedie’s trim-and-fill method (25) was applied using the R package meta (version 2024.12.0). This method estimates the number of potentially missing studies due to asymmetry in the funnel plot and imputes them to provide an adjusted effect size. In accordance with methodological guidelines, this procedure was only performed for outcomes for which Egger’s test indicated significant asymmetry. As the small number of studies within subgroups limits the reliability and interpretability of this approach, the trim-and-fill method was only applied to the overall data sets for each outcome category (26, 27).

Statistical analyses

All statistical analyses were performed using JASP version 0.19.3.0. To further investigate the effects of fatigue, subgroups were created based on performance level, distinguishing between elite and sub-elite players. In order to quantify the effects of fatigue on physiological parameters, physical fitness and stroke performance in healthy tennis players, the within-subject standardized mean difference (SMD_W) was calculated with the following formula: $SMD_W = (\text{pretest mean value} - \text{posttest mean value}) / \text{pretest standard deviation}$ (28). The SMD_W can be either positive or negative. A positive value indicates an increase in parameters, expressed by an increase in stroke performance, physical fitness or physiological parameters, from the pretest (non-fatigued) to the posttest (fatigued). Conversely, a negative value indicates a decrease in performance, expressed by a reduction in stroke performance, physical fitness or physiological parameters. SMD_W values can be classified and interpreted according to Cohen (29) into the following ranges: $0 \leq 0.49$ representing small effects, $0.50 \leq 0.79$ representing moderate effects, and ≥ 0.80 representing large effects. Further, Deeks et al. (30) postulate that heterogeneity (I^2), which reflects the proportion of variability in study results due to heterogeneity rather than random error, can be interpreted as trivial ($0 \leq 40\%$), moderate ($30 \leq 60\%$), substantial ($50 \leq 90\%$), or considerable ($75 \leq 100\%$). In cases of substantial or substantial heterogeneity ($I^2 \geq 50\%$), potential sources were explored using subgroup analyses, leave-one-out sensitivity analysis, and meta-regression models to assess the influence of performance level and study characteristics.



For the meta-analytic approach, the Sidik-Jonkman method was used due to its improved error rates in small sample scenarios and its robustness in estimating heterogeneity (31). Further, the Leave-One-Out (LOO) analysis was conducted to assess the robustness of the results and to identify potentially influential cases (32–34). This method allows for the detection of studies that disproportionately influence the overall effect

estimate, increasing the reliability of meta-analytic results. To further investigate the influence of performance level on the overall effect size, a meta-regression was performed including the subgroup as a predictor. Due to methodological differences between subgroup analyses and meta-regression models, slight variations in *SMD* estimates were observed. Specifically, while subgroup analyses estimate effect sizes independently for each

group, meta-regression considers subgroup as a covariate in a unified model, which may lead to different weighting of individual studies and slight shifts in overall effect sizes. These differences were expected and are in line with previous methodological recommendations (35). The results of both approaches are reported for transparency. To assess the robustness of the results, a sensitivity analysis was performed by excluding studies identified as influential by the leave-one-out diagnostic. The meta-analysis was repeated without these studies, and changes in effect size and heterogeneity were reported accordingly. In addition to the subgroup (performance level), we conducted exploratory meta-regressions to examine whether the age (youth vs. adults) of the participants or the type of fatigue protocol (e.g., match play, performance tests or conditioning drills) predicted the effect sizes. These additional covariates were tested in separate models to explore their potential influence on outcome variability.

Results

Study selection

Figure 1 illustrates the stages of the systematic literature search and study selection process. The initial search identified 644 articles for review and another three studies were included from other sources (e.g., reference lists, review articles). After removing duplicates and screening titles and abstracts, 46 studies were assessed for eligibility. Of these, 28 studies were excluded for the following reasons: three involved injured tennis players, five did not report relevant parameters (e.g., physiology, physical fitness and stroke performance), six lacked sufficient information on outcome measures, ten used inadequate study designs, two did not include tennis players, and two was not written in English.

Study coding

The included studies were coded in accordance with the following variables to ensure a consistent approach to data extraction: author and year of publication, number of subjects, sex, age, and the study group categorized by the type of fatigue protocol applied. To evaluate the results, three main categories of parameters were differentiated: physical fitness (e.g., countermovement jump, shuttle run), stroke performance (e.g., stroke velocity, stroke accuracy), and physiological response (e.g., blood lactate, heart rate). As some studies reported more than one variable within the same outcome category, we gave priority to the most frequently reported measure in each category to minimize the heterogeneity between studies (Table 2). For physiological measures, blood lactate was selected as the primary outcome, with creatine kinase and vital capacity as alternative measures. For physical fitness, the countermovement jump (CMJ) was the preferred outcome, while knee extension strength, center of pressure (COP) displacement, 20-m shuttle run and T-test

were considered as alternative measures. For stroke performance, serve velocity was the most reported outcome, with serve speed and serve accuracy used as alternatives when serve velocity was not available. The number of studies using each measure is displayed in Table 2.

Study characteristics

This meta-analysis includes 18 studies that investigate the impact of fatigue-inducing interventions, match play, and training on physiological parameters, physical fitness and stroke performance (see Table 3). The studies included a total of 318 tennis players, with sample sizes ranging from 6 (36) to 36 (37) participants. The age of the participants ranged from twelve to 37 years. Significant variation was observed in performance levels across the studies, with participants ranging from professional players (38) to national-level players (8, 36, 39–42), elite (37) and competitive (43) junior players, advanced-level players (9, 12, 44, 45), county players (14) and recreational players (46, 47). Furthermore, one study incorporated a combination of participants with varying degrees of expertise, including both experts and non-experts (16). Additionally, several studies involved mixed-sex cohorts (15, 16, 44, 47, 48) or did not report gender (12, 37), while others (8, 9, 36, 38–43, 45, 46) exclusively recruited male participants.

For subgroup analyses, players were classified as “elite” if they were described as professional, expert, competitive or elite, whereas “sub-elite” included recreational, county, non-expert or advanced players, based on the classifications reported in the original studies.

Outcome measures

A quantitative synthesis of the literature revealed that a total of five studies analyzed the physiological response to fatigue. Three studies focused on the analysis of blood lactate (42, 43, 46), while one study examined the role of creatine kinase (36) and vital capacity (48), respectively. A total of eleven studies were conducted to investigate the influence of fatigue on physical fitness. Five studies used the CMJ as a measure of physical fitness (36, 40, 41, 47, 48), while two studies assessed internal rotation strength (38, 44). Further, knee extension strength (37), COP displacements (45), a 20-m shuttle run (8), and the T-test

TABLE 2 Overview of the preferred and alternative outcome by category.

Category	Preferred outcome	Alternative outcome
Physiology	Blood lactate (n = 3)	Creatine kinase (n = 1) Vital capacity (n = 1)
Physical fitness	Countermovement jump (CMJ) (n = 5)	Knee extension strength (n = 1) Center of pressure displacement (n = 1) 20-m shuttle run (n = 1) T-test (n = 1)
Stroke performance	Serve velocity (n = 8)	Stroke velocity (n = 2) Stroke accuracy (n = 2)

TABLE 3 Studies examining the effects of fatigue on physiological, physical fitness, and stroke performance related parameters in healthy tennis players.

Reference	No. of participants; sex; age [years (mean ± SD or range)]; performance level	Fatigue protocol	Outcome and unit			
Vergauwen et al. (8)	20; M; 21 ± 1 years; national	Leuven Tennis Performance Test	Serve velocity [km/h] Shuttle run [s]			
Ferrauti et al. (39)	10; M; 25.3 ± 3.7 years; national	Passing shot drill with 10 s rest	Stroke velocity [km/h] Blood lactate [mmol/L]			
		Passing shot drill with 15 s rest	Stroke velocity [km/h] Blood lactate [mmol/L]			
Davey et al. (15)	18; M (9), F (9); 19–23 years; county	Loughborough Tennis Skills Test: Groundstrokes	Forehand accuracy [%]			
Maraga et al. (36)	6; M; 12.8 ± 1.2; nationally ranked	90 min single match	CMJ [cm] Stroke velocity [km/h] Creatine kinase [U/L]			
			Malliou et al. (37)	36; N/A; 14 ± 2 years; elite	90-min training session	Right knee extensors at 60°/s [Nm]
			Lyons et al. (16)	13; M (7), F (6); 19.5 ± 3.0 years, experts	Modified Loughborough Tennis Skills Test: Groundstrokes	Groundstroke accuracy [%]
11; M (13), F (4); 24.9 ± 9.6 years, non-experts	Modified Loughborough Tennis Skills Test: Groundstrokes	Groundstroke accuracy [%]				
Murphy et al. (46)	8; M; 24.2 ± 1.2 years; recreational	Cardio tennis session	Blood lactate [mmol/L]			
	8; M; 37.7 ± 6.7 years; recreational		Blood lactate [mmol/L]			
	8; M; 24.3 ± 2.6 years; recreational		Blood lactate [mmol/L]			
	8; M; 35.6 ± 2.7 years; recreational		Blood lactate [mmol/L]			
Rota et al. (12)	10; N/A; 23.8 ± 4.0 years; advanced	40-min fatiguing intermittent exercise (4 sets of intense tennis strokes)	Serve velocity [m/s]			
Gescheid et al. (40)	7; M; 21.4 ± 2.2 years; national ranking	4-h singles tennis match	Serve velocity [km/h] CMJ [cm]			
			Pialoux et al. (43)	11; M; 13.4 ± 1.3 years; competitive	Playing HIIT session	Blood lactate [mmol/L] Serve velocity [km/h]
		Non-playing HIIT session		Blood lactate [mmol/L] Serve velocity [km/h]		
Gomes et al. (41)	10; M; 16.6 ± 1.4 years; national	3-h match play	CMJ [cm]			
Martin et al. (9)	8; M; 20.4 ± 2.8 years; advanced	3-h match play	Serve velocity [m/s]			
Moreno-Perez et al. (38)	26; M; 20.4 ± 4.4; professional	Simulated tennis match (best of three)	Serve velocity [km/h] Isometric strength IR (dominant side) [N/kg]			
			Amatori et al. (47)	12; M (8), F (4); 23.0 ± 5.9 years; recreational	120-min match	CMJ [cm]
Colomar et al. (45)	15; M; 16.5 ± 1.5 years; advanced	80-min simulated match	COP displacement [mm]			
Fuentes-García et al. (48)	32; F (7), M (25); 21.4 ± 1.5 years; recreational	HIIT training	Serve velocity [km/h] CMJ [cm] Forced vital capacity [l]			
			Colomar et al. (44)	20; M (12), F (8); 16.9 ± 1.7 years; advanced	80-min simulated tennis match	Serve velocity [km/h] MVC IR [N]
				Bilic et al. (42)	21; M; 12.9 ± 0.8 years; national	300-m running test

CMJ, countermovement jump; COP, center of pressure; F, female; HIIT, high intensity interval training; IR, internal rotation, M, male; MVC, maximum voluntary contraction; NA, not available; RM, repetition maximum; SMDw, within-subject standardized mean difference.

(42) were each assessed in one study. The impact of fatigue on stroke performance in tennis was evaluated in twelve studies. Of these, eight investigated the effects of fatigue on serve speed (8, 9, 12, 38, 40, 43, 44, 48), two examined its impact on stroke speed (36, 39), and two analyzed its effects on groundstroke accuracy (15, 16).

Fatigue protocol characteristics

The included studies used different fatigue protocols. Three studies applied a tennis-specific performance test as fatigue

protocol. These were the Leuven Tennis Performance Test (8), which measures serve speed and shuttle run performance, or the Loughborough Tennis Skills Test (15, 16) and its modified versions, which assess groundstroke accuracy. In eight trials, simulated match play lasted between 40 min and 4 h (9, 36–38, 40, 41, 44, 45, 47). High intensity interval training (HIIT) tennis sessions were used in one study (43). In addition, passing shot drills with different rest intervals of 10 and 15 s were performed (39). Other protocols included a cardio tennis session (46) and a 40-min intermittent exercise protocol (12). Strength and conditioning interventions included HIIT (48) or a 300-m running test (42).

Methodological quality of the included trials

JBI critical appraisal tool

The quality assessment of the studies included revealed that all studies met ≥ 3 out of 4 criteria related to the definition and application of inclusion criteria. In addition, all studies met ≥ 2 of 3 criteria regarding the potential for bias in patient selection, and all studies met ≥ 2 of 3 criteria addressing the use of statistical methods and reporting of results (Supplementary Table S1). Overall, all included studies met at least seven out of ten criteria.

Sensitivity analysis

The LOO sensitivity analysis (Table 4 and Supplementary Table S3) was performed to assess the influence of individual studies on the heterogeneity (I^2) and SMD_w in physiology, physical fitness, and stroke performance. Exclusion of influential studies resulted in notable changes in effect sizes and heterogeneity values in the different subgroups. In the physiology category, the sub-elite subgroup showed a significant decrease in heterogeneity from considerable ($I^2 = 91.97\%$) to trivial ($I^2 = 19.82\%$) after the exclusion of an influential study (48), while the effect size increased ($SMD_w = -6.85$ to -8.46). The elite subgroup remained in the moderate heterogeneity range ($I^2 = 31.23\%$). The overall category retained substantial heterogeneity, with the I^2 decreasing slightly from 97.11% to 95.38% after exclusion. Fuentes-Garcia et al. (48) was identified as an influential study. In the physical fitness category, the elite subgroup showed the most pronounced change, with heterogeneity decreasing from substantial ($I^2 = 88.08\%$) to trivial ($I^2 = 19.29\%$), and the effect size shifting from large ($SMD_w = -0.93$) to small ($SMD_w = -0.01$). The sub-elite subgroup showed a reduction in heterogeneity from considerable ($I^2 = 91.90\%$) to substantial ($I^2 = 70.62\%$) and a slight change in effect size ($SMD_w = -0.52$ to -0.57). Across all subgroups, Bilic et al. (42) was identified as an important influential study. For stroke performance, the sub-elite subgroup showed a reduction in heterogeneity from considerable ($I^2 = 93.92\%$) to moderate ($I^2 = 38.65\%$) after exclusion, with a corresponding shift in effect size from large ($SMD_w = -0.90$) to small ($SMD_w = -0.23$). The

elite subgroup also showed a reduction in heterogeneity from substantial ($I^2 = 56.69\%$) to moderate ($I^2 = 36.69\%$). Overall, the stroke performance category showed a significant decrease in heterogeneity from substantial ($I^2 = 88.61\%$) to moderate ($I^2 = 36.38\%$). The study by Davey et al. (15) (sub-elite players) and the work by Vergauwen et al. (8) (elite players) were identified as influential.

Meta regression

Meta-regression analysis (Table 5 and Supplementary Table S4) was performed to assess the influence of subgroup classification (elite players vs. sub-elite players) on the SMD in physiology, physical fitness and stroke performance, both before and after exclusion of influential studies. In the physiology category, subgroup classification had a significant effect before exclusion ($F_{(1,8)} = 5.460$, $p = 0.048$), with the sub-elite group having a significantly lower SMD than the elite group ($\beta = -4.474$, 95% CI = -8.888 to -0.059). After exclusion, this effect became highly significant ($F_{(1,7)} = 38.155$, $p < 0.001$) and the estimated coefficient increased in magnitude ($\beta = -6.510$, 95% CI = -9.003 to -4.018), indicating a greater difference between the two groups after the removal of influential studies. For physical fitness, no significant effect of subgroup classification was observed before exclusion ($F_{(1,9)} = 0.324$, $p = 0.583$), with the subgroup coefficient ($\beta = 0.403$, 95% CI = -1.199 to 2.005) showing no meaningful differentiation between elite and sub-elite athletes. After exclusion, the effect remained non-significant ($F_{(1,7)} = 2.190$, $p = 0.182$), with a coefficient of $\beta = 0.558$, 95% CI = -0.334 to 1.451 , indicating no systematic difference between the groups before or after removal of influential studies. In stroke performance, subgroup classification was not a significant predictor of SMD before exclusion ($F_{(1,13)} = 0.606$, $p = 0.450$), with a coefficient of $\beta = -0.486$, 95% CI = -1.834 to 0.862 . After exclusion, the effect size became even smaller ($F_{(1,11)} = 0.00065$, $p = 0.980$), with the coefficient approaching zero ($\beta = 0.007$, 95% CI = -0.618 to 0.632), suggesting no differentiation between groups. In addition to performance level, exploratory meta-regression analyses were conducted to examine the influence of age group (youth vs. adult) and fatigue protocol type (match play, tennis-specific

TABLE 4 Standardized mean difference (SMD) and heterogeneity (I^2) without meta-regression.

Parameter	Group or subgroup	Before exclusion		After exclusion	
		SMD_w	I^2	SMD_w	I^2
Physiology	Sub-elite players	-6.85	91.965	-8.46	19.824
	Elite players	-1.96	31.228		
	All players	-4.12	97.107	-4.61	95.379
Physical fitness	Sub-elite players	-0.52	91.904	-0.57	70.622
	Elite players	-0.93	88.080	-0.01	19.292
	All players	-0.74	89.860	-0.31	66.549
Stroke performance	Sub-elite players	-0.90	93.923	-0.23	38.646
	Elite players	-0.39	56.691	-0.23	36.696
	All players	-0.60	88.613	-0.23	36.389

TABLE 5 Standardized mean difference (SMD) and heterogeneity (I^2) with meta-regression.

Parameter	Group or subgroup	Before exclusion		After exclusion	
		SMD_w	I^2	SMD_w	I^2
Physiology	Sub-elite players	-6.43	N/A	-8.47	N/A
	Elite players	-1.95	N/A		
	All players	-4.19	93.902	-4.85	63.022
Physical fitness	Sub-elite players	-0.52	N/A	-0.58	N/A
	Elite players	-0.93	N/A	-0.02	N/A
	All players	-0.74	90.084	-0.33	61.463
Stroke performance	Sub-elite players	-0.86	N/A	-0.23	N/A
	Elite players	-0.38	N/A	-0.23	N/A
	All players	-0.60	88.519	-0.23	37.675

NA, not available.

drills, and other protocols) on fatigue-related performance outcomes (Supplementary Table S4). While age did not significantly moderate the effects in any performance category (all $p > .05$), a significant effect of fatigue protocol on physical fitness outcomes was observed after the exclusion of influential studies ($F_{(2,6)} = 5.718$, $p = .041$). *post-hoc* analysis based on estimated marginal means indicated that tennis-specific drills were associated with the strongest fatigue-related reductions in physical fitness [SMD = -1.67, 95% CI (-2.74, -0.60)], compared to match play [SMD = -0.20, 95% CI (-0.60, 0.19)] and other protocols [SMD = 0.09, 95% CI (-0.75, 0.93)].

Egger’s test

Egger’s test revealed significant asymmetry for physiology (sub-elite players and all players, $p < 0.001$), physical fitness (sub-elite players, $p = 0.002$) and stroke performance (elite players, $p = 0.002$) subgroups. No significant asymmetry was found in the remaining subgroups (Table 6).

Trim-and-fill method

Due to the significant asymmetry detected by Egger’s test in the physiology category across all players ($p < .001$), a trim-and-fill analysis was performed to adjust for potential publication bias. The unadjusted effect size was SMD = -4.11 [95% CI: (-6.34, -1.88)], with one potentially missing study imputed. After adjustment, the effect size decreased to SMD = -3.54 [95% CI: (-5.93, -1.15)], suggesting a slight overestimation in the original estimate. The corresponding funnel plot is shown in Supplementary Figure S5.

Effects of fatigue on physiological measures

The impact of fatigue on physiological parameters was evaluated through meta-regression analysis (Table 5). As demonstrated in Figure 2a, the sub-elite group showed a large effect (SMD_w = -6.43) prior to exclusion. After exclusion (Figure 2b), the effect size underwent a further increase (SMD_w = -8.47), indicating still a large effect of fatigue. For the elite group, the effect size remained stable before and after exclusion (SMD_w = -1.95 vs. -1.96), both indicating large effects. The initial evaluation of the overall physiology category revealed considerable heterogeneity ($I^2 = 93.90%$), which subsequently

diminished to a substantial level ($I^2 = 63.02%$), accompanied by an augmentation in effect size from SMD_w = -4.19 to -4.85 following exclusion.

Effects of fatigue on measures of physical fitness

The impact of fatigue on physical fitness measures was analyzed through meta-regression (Table 5). Prior to the exclusion of data (Figure 3a), the elite group demonstrated a moderate effect (SMD_w = -0.93). After the exclusion of data (Figure 3b), the effect size underwent a substantial shift to small (SMD_w = -0.01). For the sub-elite group, the effect size was moderate before exclusion (SMD_w = -0.52) and small after exclusion (SMD_w = -0.33). The overall physical fitness category initially demonstrated considerable heterogeneity ($I^2 = 90.08%$), which decreased to substantial levels ($I^2 = 61.46%$), while the effect size shifted from moderate (SMD_w = -0.74) to small (SMD_w = -0.31).

Effects of fatigue on measures of stroke performance

The impact of fatigue on stroke performance was examined using meta-regression (Table 5). Before the exclusion of data, the sub-elite group exhibited a large effect (SMD_w = -0.86). After data exclusion, the effect size changed to SMD_w = -0.23, indicating a small effect (Figures 4a,b). For the elite group, the effect size was small before exclusion (SMD_w = -0.38) and remained small after exclusion (SMD_w = -0.23). The stroke performance category exhibited substantial heterogeneity ($I^2 = 88.52%$) at the outset, which diminished to moderate level ($I^2 = 37.67%$) following exclusion. The effect size changed from moderate (SMD_w = -0.60) to small (SMD_w = -0.23).

Discussion


The objective of this systematic review and meta-analysis was to investigate the effects of fatigue protocols on physiological, physical

TABLE 6 Results for the eggers’ test used to assess publication bias.

Parameter	Group or subgroup	z-value	95% CI	p-value	Eggers’ test ^a
Physiology	Sub-elite players	-9.636	0.773 to 2.540	<.001	Asymmetry
	Elite players	0.255	-13.218 to 6.709	.799	No asymmetry
	All players	-8.524	0.573 to 2.600	<.001	Asymmetry
Physical fitness	Sub-elite	-3.096	0.528 to 4.046	.002	Asymmetry
	Elite players	0.001	-4.146 to 2.292	.999	No asymmetry
	All players	-1.551	-1.217 to 2.721	.121	No asymmetry
Stroke performance	Sub-elite players	0.236	-5.889 to 3.209	.813	No asymmetry
	Elite players	-3.060	0.190 to 3.162	.002	Asymmetry
	All players	-1.476	-1.323 to 2.830	.140	No asymmetry

95% CI, 95% confidence interval.

^aIndicates the presence of publication bias.




fitness, and stroke performance parameters in healthy tennis players. The results demonstrate that fatigue has large negative effects on physiological parameters, moderate effects on physical fitness, and small effects on stroke performance. Furthermore, subgroup analyses revealed that elite players exhibited greater resistance to fatigue compared to sub-elite players, particularly in physiological and physical fitness measures. The exclusion of influential studies resulted in the enhancement of effect sizes in physiology and a substantial reduction in heterogeneity, thereby confirming the robustness of the observed effects.

Fatigue protocols revealed a large effect on physiological parameters with higher blood lactate levels, heart rate, and creatine concentrations. These effects, as well as metabolic disturbances, are also highlighted in studies describing fatigue as a key mechanism for performance decline in tennis (19, 20). The metabolic and thermal stress that threatens physiological stability


is particularly elevated during long matches (10). Gescheit et al. (40) showed increasing levels of creatine kinase during 4-h matches for four consecutive days, indicating an accumulation of physiological damage.

Prior to the exclusion of influential studies, the sub-elite group demonstrated heightened fatigue-induced physiological responses in comparison to the elite group, exhibiting an effect size of $SMD_w = -6.43$. After the exclusion, the effect size underwent a further augmentation ($SMD_w = -8.47$), thereby indicating that the initial analysis may have underestimated the impact of fatigue in this subgroup. Furthermore, heterogeneity decreased from considerable to substantial, thereby supporting the robustness of the findings. The increase in effect size due to the exclusion of studies suggests that some studies may have underestimated the effects of fatigue due to methodological differences, such as shorter protocol duration or reduced training



intensity of the participants. Fuentes-Garcia et al. (48) assessed forced vital capacity whereas Murphy et al. (46) measured blood lactate levels. This methodological discrepancy may have contributed to the exclusion of the Fuentes-Garcia et al. (48) study, as blood lactate is a more commonly used marker for fatigue assessment in physiological analyses. Meta-regression confirmed that subgroup classification (elite players vs. sub-elite players) significantly influenced physiological effects before exclusion and became even more pronounced after exclusion. These findings imply that higher-trained athletes may possess superior physiological adaptations that enhance fatigue resistance, as previously hypothesized in other studies (16). While fatigue primarily affected physiological responses, its impact on physical fitness parameters was more moderate, with notable variations

between elite and sub-elite tennis players, with performance deteriorations being observed in agility time, countermovement jump height, and sprint time. Initially, the elite group exhibited a large effect size ($SMD_w = -0.93$), which declined to small ($SMD_w = -0.01$) after exclusion. Following the exclusion of studies, the overall SMD_w was reduced from -0.74 to -0.31 , indicating that the effects were small. This finding suggests that influential studies may have overestimated the fatigue-induced performance impairments in the elite subgroup. A possible explanation for this is that Bilic et al. (42) was the only study in this category and subgroup that did not implement a tennis-specific intervention, contributing to methodological differences. This difference in study design may have influenced the estimated effect size, further justifying its exclusion from the



sensitivity analysis. For the sub-elite group, effect sizes remained relatively stable before ($SMD_w = -0.52$, moderate effect) and after study exclusion ($SMD_w = -0.57$, moderate effect). In the overall physical fitness category, heterogeneity was initially considerable ($I^2 = 90.08\%$) but decreased to substantial levels ($I^2 = 66.55\%$) after exclusion. Meta-regression analysis revealed no significant subgroup differences before or after exclusion. These findings indicate that physical fitness parameters are influenced by fatigue, but elite players may mitigate these effects more efficiently through neuromuscular and cardiovascular adaptations (49).

Compared to physiological and physical fitness measures, stroke performance appeared to be the most resilient to fatigue effects

($SMD_w = -0.60$), indicating that stroke performance is the most robust parameter of the three outcomes investigated in relation to fatigue, as it is dependent on motor learning (50). However, Rota et al. (12) reported biomechanical adaptations resulting in a reduction in stroke performance and accuracy. In contrast, professional tennis players are able to maintain serve speed over five sets (51). Prior to the implementation of exclusion, the sub-elite group demonstrated a substantial effect ($SMD_w = -0.86$), which diminished to a small effect ($SMD_w = -0.23$) after the exclusion process. In a similar manner, the elite group initially exhibited a moderate effect ($SMD_w = -0.38$), which remained consistent following the exclusion procedure ($SMD_w = -0.23$). The stroke

performance category exhibited considerable heterogeneity ($I^2 = 88.52\%$) at the outset, which diminished to moderate levels ($I^2 = 37.68\%$) following the exclusion of outlying values. Overall SMD_w was reduced due to study exclusion to a small effect ($SMD_w = -0.23$). The reduction in heterogeneity after exclusion indicates that some studies introduced variability, potentially due to different fatigue assessment methods or player' competition levels. Meta-regression analysis revealed that subgroup classification did not serve as a significant predictor, both before and after the exclusion process. This finding indicates that the effects of fatigue on stroke performance remain relatively stable across a range of competition levels.

Our results showed that physiological parameters ($SMD_w = -4.19$) were most negatively affected by fatigue, followed by physical fitness parameters ($SMD_w = -0.74$), and stroke performance related parameters ($SMD_w = -0.60$). Therefore, physiological recovery should be targeted by specific measures. This can be done through hydration or nutrition during competition (52, 53). Additionally, research indicates that nutritional strategies, including carbohydrate supplementation, can help reduce performance declines caused by fatigue (54). Post-exercise strategies should then be used to promote lactate clearance and accelerate muscle recovery. Studies have shown that strategies such as cold baths, active recovery, and compression garments can aid in this process (55–57). In order to minimize fatigue-effects on physical fitness, high-intensity interval training may be advisable (58). Plyometric training can also improve neuromuscular efficiency, reducing the physiological cost of rapid changes in direction (59). For skill training, care should be taken to ensure that new skills are learned in a non-fatiguing state (60). Specifically, Davey et al. (15) recommend avoiding lactate concentrations >8 mmol/L, heart rates >180 bpm, and perceived exertion >16 during technique training to stabilize tennis skills. On the other hand, technique training under fatigue conditions can be useful in order to maintain biomechanical efficiency under load and to keep stroke speed and accuracy stable over a longer period of time (12, 61). The various differences in performance during training and recovery should also be taken into account. Elite players, who typically have higher neuromuscular efficiency and autonomic recovery capacity (62), benefit from precision recovery protocols (e.g., whole-body cryotherapy, active recovery, compression garments) to maintain high training frequency and manage accumulated load (63). In contrast, sub-elite athletes often show greater declines in performance with fatigue and may require fundamental improvements in anaerobic and strength capacity through high-intensity interval training (HIIT), plyometrics, and neuromuscular coordination work (64). In addition, individualized periodization models—particularly for sub-elite athletes—should ensure sufficient recovery time and controlled exposure to fatigue to support technical learning without performance collapse (65).

Limitations

The present meta-analysis has several limitations. First, the number of studies per category is relatively small ($n = 5–12$), which limits the statistical power and robustness of the findings. Future

replication studies with larger samples are needed to confirm and generalize the observed effects. Secondly, methodological differences between fatigue protocols (e.g., match play vs. local fatigue) make direct comparisons difficult. While local fatigue protocols offer high internal validity by minimizing external variables, they often lack ecological validity. In contrast, match play protocols better reflect real game conditions but introduce uncontrolled factors. An integrated approach, such as replicating match-play situations in controlled laboratory settings (e.g., using virtual reality), may help to bridge this gap. Third, heterogeneity in participant characteristics (e.g., gender, age, and skill level) increases the variability of results. Due to limited data, it was not possible to differentiate by gender or age; instead, subgroup analyses based on performance level (elite vs. sub-elite) were more appropriate. Fourth, Egger's test indicated potential publication bias and small study effects, particularly for physiological and physical fitness outcomes. Future studies could mitigate this by pre-registering protocols and increasing sample sizes to ensure more balanced evidence. Furthermore, as all included studies focused on short-term responses (<24 h), evidence on long-term or chronic fatigue adaptations remains unexplored—highlighting the need for longitudinal designs. Finally, the under-representation of female athletes limits the ability to draw gender-specific conclusions. Given the known sex differences in fatigue resistance and recovery patterns, future research should explicitly address female-specific responses in tennis-related fatigue.

Conclusions

This meta-analysis confirms that fatigue exerts the most significant effects on physiological parameters, followed by physical fitness and stroke performance. The exclusion of influential studies resulted in effect sizes that were more pronounced and reduced heterogeneity, suggesting that some studies may have overestimated the fatigue effects, particularly in the sub-elite group. In contrast, elite players demonstrated greater resistance to fatigue, especially in physiological and physical fitness measures, thereby supporting the notion that training adaptations play a crucial role in fatigue management. Future research should explore longitudinal training interventions to enhance fatigue resistance, particularly in sub-elite tennis players, and investigate optimal recovery protocols for physiological stabilization in competitive settings.

Data availability statement

The original contributions presented in the study are included in the article/[Supplementary Material](#), further inquiries can be directed to the corresponding author.

Author contributions

JL: Conceptualization, Data curation, Formal analysis, Methodology, Writing – original draft, Writing – review &

editing, TM: Conceptualization, Data curation, Writing – review & editing.

Funding

The author(s) declare that financial support was received for the research and/or publication of this article. Support was provided through the Open Access Publication Fund of the University of Duisburg-Essen, organised by the project DEAL. The funding body is independent of the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Chalder T, Cella M. Fatigue. In: Davey P, Sprigings D, editors. *Diagnosis and Treatment in Internal Medicine (online ed., Chapter 79)*. Oxford: Oxford University Press (2018). doi: 10.1093/med/9780199568741.003.0079
- Enoka RM, Duchateau J. Muscle fatigue: what, why and how it influences muscle function. *J Physiol.* (2008) 586:11–23. doi: 10.1113/jphysiol.2007.139477
- Taylor JL, Amann M, Duchateau J, Meeusen R, Rice CL. Neural contributions to muscle fatigue: from the brain to the muscle and back again. *Med Sci Sports Exerc.* (2016) 48:2294–306. doi: 10.1249/MSS.0000000000000923
- Behrens M, Gube M, Chaabene H, Prieske O, Zenon A, Broscheid K-C, et al. Fatigue and human performance: an updated framework. *Sports Med.* (2023) 53:7–31. doi: 10.1007/s40279-022-01748-2
- Enoka RM, Duchateau J. Translating fatigue to human performance. *Med Sci Sports Exerc.* (2016) 48:2228–38. doi: 10.1249/MSS.0000000000000929
- Amann M. Central and peripheral fatigue: interaction during cycling exercise in humans. *Med Sci Sports Exerc.* (2011) 43:2039–45. doi: 10.1249/MSS.0b013e31821f59ab
- Meeusen R, Watson P, Hasegawa H, Roelands B, Piacentini MF. Central fatigue: the serotonin hypothesis and beyond. *Sports Med.* (2006) 36:881–909. doi: 10.2165/00007256-200636100-00006
- Vergauwen L, Spaepen AJ, Lefevre J, Hespel P. Evaluation of stroke performance in tennis. *Med Sci Sports Exerc.* (1998) 30:1281–8. doi: 10.1097/00005768-199808000-00016
- Martin C, Bideau B, Delamarche P, Kulpa R. Influence of a prolonged tennis match play on serve biomechanics. *PLoS One.* (2016) 11:e0159979. doi: 10.1371/journal.pone.0159979
- Mendez-Villanueva A, Fernandez-Fernandez J, Bishop D. Exercise-induced homeostatic perturbations provoked by singles tennis match play with reference to development of fatigue. *Br J Sports Med.* (2007) 41:717–22; discussion 722. doi: 10.1136/bjism.2007.037259
- Hornery DJ, Farrow D, Mujika I, Young W. An integrated physiological and performance profile of professional tennis. *BJSM.* (2007) 41:531–6; discussion 536. doi: 10.1136/bjism.2006.031351
- Rota S, Morel B, Saboul D, Rogowski I, Hautier C. Influence of fatigue on upper limb muscle activity and performance in tennis. *J Electromyogr Kinesiol.* (2014) 24:90–7. doi: 10.1016/j.jelekin.2013.10.007
- Fernandez-Fernandez J, Moya-Ramon M, Santos-Rosa FJ, Gantois P, Nakamura FY, Sanz-Rivas D, et al. Within-session sequence of the tennis serve training in youth elite players. *Int J Environ Res Public Health.* (2020) 18:244. doi: 10.3390/ijerph18010244
- Davey PR, Thorpe RD, Williams C. Simulated tennis matchplay in a controlled environment. *J Sports Sci.* (2003) 21:459–67. doi: 10.1080/0264041031000101926

Generative AI statement

The author(s) declare that no Generative AI was used in the creation of this manuscript.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fspor.2025.1578914/full#supplementary-material>

- Davey PR, Thorpe RD, Williams C. Fatigue decreases skilled tennis performance. *J Sports Sci.* (2002) 20:311–8. doi: 10.1080/026404102753576080
- Lyons M, Al-Nakeeb Y, Hankey J, Nevill A. The effect of moderate and high-intensity fatigue on groundstroke accuracy in expert and non-expert tennis players. *J Sports Sci Med.* (2013) 12:298–308.
- Turner M, Beranek P, Sahrom S, Lo J, Ferrauti A, Dunican IC, et al. The impact of sleep behaviours, chronotype and time of match on the internal and external outcomes of a tennis match. *Int J Sports Sci Coach.* (2023) 18:2099–107. doi: 10.1177/17479541221130443
- Girard O. Neuromuscular fatigue in tennis: mind over muscle? *Int Tennis Federat.* (2014) 63:10–3. doi: 10.52383/itfcoaching.v22i63.499
- Reid M, Duffield R. The development of fatigue during match-play tennis. *Br J Sports Med.* (2014) 48(Suppl 1):i7–11. doi: 10.1136/bjsports-2013-093196
- Hornery DJ, Farrow D, Mujika I, Young W. Fatigue in tennis: mechanisms of fatigue and effect on performance. *Sports Med.* (2007) 37:199–212. doi: 10.2165/00007256-200737030-00002
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med.* (2009) 151:264–9; W64. doi: 10.7326/0003-4819-151-4-200908180-00135
- Girard O, Christian RJ, Racinais S, Périard JD. Heat stress does not exacerbate tennis-induced alterations in physical performance. *Br J Sports Med.* (2014) 48(Suppl 1):i39–44. doi: 10.1136/bjsports-2013-093165
- Munn Z, Barker TH, Moola S, Tufanaru C, Stern C, McArthur A, et al. Methodological quality of case series studies: an introduction to the JBI critical appraisal tool. *JBI Evid Synth.* (2020) 18:2127–33. doi: 10.11124/JBISIR-D-19-00099
- Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *Br Med J.* (1997) 315:629–34. doi: 10.1136/bmj.315.7109.629
- Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics.* (2000) 56:455–63. doi: 10.1111/j.0006-341x.2000.00455.x
- Peters JL, Sutton AJ, Jones DR, Abrams KR, Rushton L. Performance of the trim and fill method in the presence of publication bias and between-study heterogeneity. *Stat Med.* (2007) 26:4544–62. doi: 10.1002/sim.2889
- Shi L, Lin L. The trim-and-fill method for publication bias: practical guidelines and recommendations based on a large database of meta-analyses. *Medicine (Baltimore).* (2019) 98:e15987. doi: 10.1097/MD.00000000000015987
- Deeks JJ, Higgins JP. Statistical algorithms in review manager 5 (2010).

29. Cohen J. Statistical power analysis. *Curr Dir Psychol Sci.* (1992) 1:98–101. doi: 10.1111/1467-8721.ep10768783
30. Deeks JJ, Higgins JP, Altman DG. Analysing data and undertaking meta-analyses. In: Higgins J, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions.* Hoboken, NJ, Chichester: Wiley Blackwell (2019), p. 241–84.
31. IntHout J, Ioannidis JP, Borm GF. The hartung-knapp-sidik-jonkman method for random effects meta-analysis is straightforward and considerably outperforms the standard DerSimonian-laird method. *BMC Med Res Methodol.* (2014) 14:25. doi: 10.1186/1471-2288-14-25
32. Harrer M, Cuijpers P, Furukawa TA, Ebert DD. *Doing Meta-analysis with R: A Hands-on Guide.* Boca Raton, London, New York: CRC Press Taylor & Francis Group (2022), p. 474 p.
33. Viechtbauer W, Cheung MW-L. Outlier and influence diagnostics for meta-analysis. *Res Synth Methods.* (2010) 1:112–25. doi: 10.1002/jrsm.11
34. Meng Z, Wang J, Lin L, Wu C. Sensitivity analysis with iterative outlier detection for systematic reviews and meta-analyses. *Stat Med.* (2024) 43:1549–63. doi: 10.1002/sim.10008
35. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J. Stat. Soft.* (2010) 36:1–48. doi: 10.18637/jss.v036.i03
36. Maraga N, Duffield R, Gescheit D, Perri T, Reid M. Playing not once, not twice but three times in a day: the effect of fatigue on performance in junior tennis players. *Int J Perform Anal Sport.* (2018) 18:104–14. doi: 10.1080/24748668.2018.1452110
37. Malliou VJ, Beneka AG, Gioftsidou AF, Malliou PK, Kallistratos E, Pafis GK, et al. Young tennis players and balance performance. *J. Strength Cond. Res.* (2010) 24:389–93. doi: 10.1519/JSC.0b013e3181c068f0
38. Moreno-Pérez V, López-Samanes Á, Domínguez R, Fernández-Eliás VE, González-Frutos P, Fernández-Ruiz V, et al. Acute effects of a single tennis match on passive shoulder rotation range of motion, isometric strength and serve speed in professional tennis players. *PLoS One.* (2019) 14:e0215015. doi: 10.1371/journal.pone.0215015
39. Ferrauti A, Pluim BM, Weber K. The effect of recovery duration on running speed and stroke quality during intermittent training drills in elite tennis players. *J. Sports Sci.* (2001) 19:235–42. doi: 10.1080/026404101750158277
40. Gescheit DT, Cormack SJ, Reid M, Duffield R. Consecutive days of prolonged tennis match play: performance, physical, and perceptual responses in trained players. *Int J Sports Physiol Perform.* (2015) 10:913–20. doi: 10.1123/ijspp.2014-0329
41. Gomes RV, Cunha VC, Zourdos MC, Aoki MS, Moreira A, Fernandez-Fernandez J, et al. Physiological responses of young tennis players to training drills and simulated match play. *J. Strength Cond. Res.* (2016) 30:851–8. doi: 10.1519/JSC.0000000000001159
42. Bilić Z, Sinković F, Barbaros P, Novak D, Zemkova E. Exercise-induced fatigue impairs change of direction performance and serve precision among young male tennis players. *Sports (Basel).* (2023):11:111. doi: 10.3390/sports11060111
43. Pialoux V, Genevois C, Capoen A, Forbes SC, Thomas J, Rogowski I. Playing vs. nonplaying aerobic training in tennis: physiological and performance outcomes. *PLoS One.* (2015) 10:e0122718. doi: 10.1371/journal.pone.0122718
44. Colomar J, Corbi F, Baiget E. Force-time curve variable outcomes following a simulated tennis match in junior players. *J Sports Sci Med.* (2022) 21:245–52. doi: 10.52082/jssm.2022.245
45. Colomar J, Corbi F, Baiget E. Alterations in mechanical muscle characteristics and postural control induced by tennis match-play in young players. *PeerJ.* (2021) 9:e11445. doi: 10.7717/peerj.11445
46. Murphy AP, Duffield R, Kellett A, Reid M. A comparison of the perceptual and technical demands of tennis training, simulated match play, and competitive tournaments. *Int J Sports Physiol Perform.* (2016) 11:40–7. doi: 10.1123/ijspp.2014-0464
47. Amatori S, Gobbi E, Moriondo G, Gervasi M, Sisti D, Rocchi MB, et al. Effects of a tennis match on perceived fatigue, jump and sprint performances on recreational players. *Open Sports Sci J.* (2020) 13:54–9. doi: 10.2174/1875399X02013010054
48. Fuentes-García JP, Díaz-García J, López-Gajardo MÁ, Clemente-Suarez VJ. Effects of combined HIIT and stroop on strength manifestations, serve speed and accuracy in recreational tennis players. *Sustainability.* (2021) 13:7717. doi: 10.3390/SU13147717
49. Johnston MJ, Cook CJ, Drake D, Costley L, Johnston JP, Kilduff LP. The neuromuscular, biochemical, and endocrine responses to a single-session vs. double-session training day in elite athletes. *J Strength Cond Res.* (2016) 30:3098–106. doi: 10.1519/JSC.0000000000001423
50. Wolpert DM, Flanagan JR. Motor learning. *Curr Biol.* (2010) 20:R467–72. doi: 10.1016/j.cub.2010.04.035
51. Maquiritiaín J, Baglione R, Cardey M. Male professional tennis players maintain constant serve speed and accuracy over long matches on grass courts. *Eur J Sport Sci.* (2016) 16:845–9. doi: 10.1080/17461391.2016.1156163
52. Robson-Ansley PJ, Gleeson M, Ansley L. Fatigue management in the preparation of Olympic athletes. *J Sports Sci.* (2009) 27:1409–20. doi: 10.1080/02640410802702186
53. Ranchordas M, Rogerson D, Ruddock A, Killer S, Winter E. Nutrition for tennis: practical recommendations. *J. Sports Sci. Med.* (2013) 12(2):211–24.
54. Brink-Elfegoun T, Ratel S, Leprêtre P-M, Metz L, Ennequin G, Doré E, et al. Effects of sports drinks on the maintenance of physical performance during 3 tennis matches: a randomized controlled study. *J Int Soc Sports Nutr.* (2014) 11:46. doi: 10.1186/s12970-014-0046-7
55. Kovacs MS, Baker LB. Recovery interventions and strategies for improved tennis performance. *Br J Sports Med.* (2014) 48(Suppl 1):i18–21. doi: 10.1136/bjsports-2013-093223
56. Fleming JA, Naughton RJ, Harper LD. Investigating the nutritional and recovery habits of tennis players. *Nutrients.* (2018) 10:443. doi: 10.3390/nu10040443
57. Karadavut Ö, Acar G. Effects of sports massage on post-workout fatigue. *Uluslararası Türk Spor ve Egzersiz Psikolojisi Dergisi.* (2024) 4:15–29. doi: 10.55376/ijtssep.1486062
58. O'Leary TJ, Collett J, Howells K, Morris MG. Endurance capacity and neuromuscular fatigue following high- vs moderate-intensity endurance training: a randomized trial. *Scand J Med Sci Sports.* (2017) 27:1648–61. doi: 10.1111/sms.12854
59. Asadi A, Arazi H, Young WB, Sáez de Villarreal E. The effects of plyometric training on change-of-direction ability: a meta-analysis. *Int J Sports Physiol Perform.* (2016) 11:563–73. doi: 10.1123/ijspp.2015-0694
60. Suna G, Alp M, Çetinkaya E. Investigating the effects of technical trainings applied 10–12 age male tennis players on their stroke performances. *HumanSciences.* (2016) 13:4980. doi: 10.14687/JHS.V13I3.4019
61. Brechbuhl C, Girard O, Millet GP, Schmitt L. Differences within elite female tennis players during an incremental field test. *Med Sci Sports Exercise.* (2018) 50:2465–73. doi: 10.1249/MSS.0000000000001714
62. Djokic Z, Zagatto AM. Ultra-short-term heart rate recovery after maximal exercise in elite European table tennis players. *Sport Sci Health.* (2019) 15:343–50. doi: 10.1007/s11332-018-0520-8
63. Poignard M, Guilhem G, de Laroche Lambert Q, Montalvan B, Bieuzen F. The impact of recovery practices adopted by professional tennis players on fatigue markers according to training type clusters. *Front Sports Act Living.* (2020) 2:109. doi: 10.3389/fspor.2020.00109
64. Lambrich J, Muehlbauer T. Physical fitness and stroke performance in healthy tennis players with different competition levels: a systematic review and meta-analysis. *PLoS One.* (2022) 17:e0269516. doi: 10.1371/journal.pone.0269516
65. Kerr AD. The physiological demands of tennis and how to train them: a case study of a sub elite female tennis player. *J Aust Strength Cond.* (2015) 23(1):25–37.

DuEPublico

Duisburg-Essen Publications online

UNIVERSITÄT
DUISBURG
ESSEN

Offen im Denken

ub | universitäts
bibliothek

This text is made available via DuEPublico, the institutional repository of the University of Duisburg-Essen. This version may eventually differ from another version distributed by a commercial publisher.

DOI: 10.3389/fspor.2025.1578914

URN: urn:nbn:de:hbz:465-20250918-102952-9



This work may be used under a Creative Commons Attribution 4.0 License (CC BY 4.0).